



UNIVERSITY OF  
LIVERPOOL

Management  
School

**Working Paper in Economics**

**# 202207**

# **Testing Granger Non-Causality in Expectiles**

**Taoufik Bouezmarni  
Mohamed Doukali  
Abderrahim Taamouti**

# Testing Granger non-causality in Expectiles

Taoufik Bouezmarni\*    Mohamed Doukali†    Abderrahim Taamouti‡

March 7, 2022

## ABSTRACT

This paper aims to derive a consistent test of Granger causality at a given expectile. We also propose a sup-Wald test for jointly testing Granger causality at all expectiles that has the correct asymptotic size and power properties. Expectiles have the advantage of capturing similar information as quantiles, but they also have the merit of being much more straightforward to use than quantiles, since they are defined as least squares analogue of quantiles. Studying Granger causality in expectiles is practically simpler and allows us to examine the causality at all levels of the conditional distribution. Moreover, testing Granger causality at all expectiles provides a sufficient condition for testing Granger causality in distribution. A Monte Carlo simulation study reveals that our tests have good finite-sample size and power properties for a variety of data-generating processes and different sample sizes. Finally, we provide two empirical applications to illustrate the usefulness of the proposed tests.

**Keywords:** Granger causality in expectiles, Granger causality in distribution, expectile regression function, asymmetric loss function, sup-Wald test.

**Journal of Economic Literature classification:** C12, C22

---

\*Département de Mathématiques, Université de Sherbrooke, CIREQ, CREAS. Email addresses: Taoufik.Bouezmarni@USherbrooke.ca

†McGill University and Centre Interuniversitaire de Recherche en Analyse des Organisations (CIRANO). Department of Economics, McGill University, Leacock Building, 855 Sherbrooke Street West, Montreal, Quebec H3A 2T7, Canada. Email: mohamed.doukali@mail.mcgill.ca

‡ *Corresponding author.* Department of Economics, University of Liverpool Management School. Address: Chatham St, Liverpool L69 7ZH. E-mail: Abderrahim.Taamouti@liverpool.ac.uk.

# Testing Granger non-causality in Expectiles

## ABSTRACT

This paper aims to derive a consistent test of Granger causality at a given expectile. We also propose a sup-Wald test for jointly testing Granger causality at all expectiles that has the correct asymptotic size and power properties. Expectiles have the advantage of capturing similar information as quantiles, but they also have the merit of being much more straightforward to use than quantiles, since they are defined as least squares analogue of quantiles. Studying Granger causality in expectiles is practically simpler and allows us to examine the causality at all levels of the conditional distribution. Moreover, testing Granger causality at all expectiles provides a sufficient condition for testing Granger causality in distribution. A Monte Carlo simulation study reveals that our tests have good finite-sample size and power properties for a variety of data-generating processes and different sample sizes. Finally, we provide two empirical applications to illustrate the usefulness of the proposed tests.

**Keywords:** Granger causality in expectiles, Granger causality in distribution, expectile regression function, asymmetric loss function, sup-Wald test.

**Journal of Economic Literature classification:** C12, C22

# 1 Introduction

Causal relations between economic and/or financial time series are typically examined using the popular concept of Granger causality introduced by Wiener (1956) and Granger (1969). While the concept is naturally defined in terms of conditional distribution [see Granger (1980) and Granger and Newbold (1986)], early studies often focus on the conditional mean [see e.g. Granger (1969)] or the conditional variance [see e.g. Granger et al. (1986) and Cheung and Ng (1996)]. Causality in other aspects of the conditional distribution of the variables of interest (e.g. high-order conditional moments, quantiles/expectiles) has been less studied in practice, despite empirical evidence showing that for many economic and financial time series, e.g. returns and output, conditional quantiles/expectiles are predictable, but not the conditional mean; see Taylor, J. W. (2008), Lee and Yang (2012), Cenesizoglu and Timmermann (2008), and Chuang (2009) among others. In this paper, we pay particular attention to the concept of expectiles introduced by Newey and Powell (1987), which provides a more complete picture of the conditional distribution of variable interest. In particular, we define Granger non-causality in expectiles and propose a parametric test for it. When all expectiles are considered jointly, the proposed test is equivalent to testing Granger non-causality in distribution. Rather than checking a necessary condition for Granger non-causality, our approach analyzes a continuous space of conditional expectile functions that fully characterizes non-causality in distribution.

The theory of Wiener-Granger causality has generated a considerable literature; for reviews, see Lütkepohl (2005), Boudjellaba, Dufour and Roy (1992, 1994), and Dufour and Taamouti (2010) and the references therein. However, to the best of our knowledge, no test has been proposed to test Granger non-causality in expectiles. Most of the existing tests focus on Granger causality in mean, thus they cannot be used in the presence of causality in expectiles. Several tests and measures have been proposed to detect Granger causality in distribution; see Su and White (2007, 2008), Taamouti et al. (2014), Bouezmarni and Taamouti (2014), Bouezmarni et al. (2012) and the references therein. However, these tests and measures are not informative about the level(s) (mean, variance, other high-order moments, expectiles) of distribution where the causality exists. To overcome this issue, Song and Taamouti (2020) consider measures of Granger causality in quantiles; see also Troster (2018).

Another interesting way of looking at causality across different levels of conditional distri-

bution, particularly the tails, is by using the concept of expectile. The latter dates back to Newey and Powell (1987), but recently re-gained a lot interest in both theoretical and applied work in econometrics and finance, see Taylor (2008), Sobotka and Kneib (2012), Holzmann and Klar (2016), Kratschmer and Zahle (2017), and Daouia et al. (2019) among others. Expectiles have the advantage of capturing similar information as quantiles, but they also have the merit of being much more straightforward to use than quantiles, since they are defined as least squares analogue of quantiles. Studying Granger causality in expectiles is practically simpler and can help us examine the causality in distribution. Roughly speaking, expectiles can be seen as a mixture of mean and quantiles. On the one hand, they are determined by an asymmetrically weighted deviations criterion, where the  $L_1$ -norm of quantiles is replaced by the  $L_2$ -norm of expectation. On the other hand, they represent a generalization of the ordinary least square regression - 50% expectile regression is the classical mean regression. In addition, using expectiles instead of quantiles has the following advantages. First, expectiles lie in the computational expedience. Indeed, the estimation based on expectile regression reduces to weighted least squares fits since the optimality criterion is differentiable with respect to the regression effects, while linear programming routines have to be used in case of quantile regression. The second advantage is that inference on expectiles is much easier than inference on quantiles, and their estimation makes more efficient use of the available data since weighted least squares rely on the distance to data points, while empirical quantiles only utilize the information on whether an observation is below or above the predictor; see Newey and Powell (1987), Abdous and Remillard (1995), and Sobotka and Kneib (2012) among others. Finally, as a risk measure it has been shown that expectiles have the attractive property of coherence [see e.g. Bellini et al. (2014)], while quantiles suffer from the lack of subadditivity. Expectiles were shown to be the only coherent and elicitable risk measures in Ziegel (2016). Further discussion and application of expectiles as risk measures are given in Taylor (2008), Kuan et al. (2009), Emmer et al. (2015), Delbaen (2013), Bellini and Di Bernardino (2017), and Daouia et al. (2018).

In this paper, we propose a consistent parametric test of Granger causality in expectiles, which is based on expectile regressions. We first propose a Wald-type test for non-causality at a fixed  $\tau$ -expectile, for  $\tau \in (0, 1)$ . Thereafter, we propose a sup-Wald test for testing Granger causality at all expectiles. The latter test checks for the joint statistical significance of the parameters of expectile regressions for all  $\tau$  in  $(0, 1)$ , which is consistent against any deviation

from non-causality in distribution, as opposed to the conventional tests of non-causality in moments. Testing Granger non-causality at all expectiles provides a sufficient condition for testing Granger non-causality in distribution. The proposed sup-Wald test statistic has the correct asymptotic size and power properties. A Monte Carlo simulation study reveals that our tests have good finite-sample size and power properties for a variety of data-generating processes and different sample sizes. We also compare the properties (size and power) of our *expectile-based* test with the existing *quantile-based* test for different values of  $\tau$ , see e.g. Koenker and Machado (1999). The simulation results show that our test outperforms the test based on quantile regression even for weak degree of causality and for both small and large samples. Finally, we provide two empirical applications to illustrate the usefulness of our tests. In these applications, we re-examine the Granger causality from volume and exchange rates to stock returns using expectile.

The rest of this paper is organized as follows. In Section 2, we introduce the notations and define the concept of Granger non-causality in expectiles. In Section 3, we consider a Wald-type test and a sup-Wald test for testing Granger causality at a given expectile and all expectiles jointly, respectively. In Section 4, we use Monte Carlo simulations to investigate the finite sample size and power properties of our tests, and we compare their performance to those of quantile-based test. Section 5 contains two applications using economic and financial data and Section 6 concludes. The proofs of the theoretical results can be found in the Appendix.

## 2 Granger causality in expectiles

We consider two variables of interest  $Y$  and  $Z$ . Let  $\{(Y_t, Z_t) : t \in \mathbb{Z}\}$  be the strictly stationary and ergodic time series process of  $Y$  and  $Z$  defined on some probability space  $(\Omega, F, P)$ . We denote  $L^2 \equiv L^2(\Omega, F, P)$  a Hilbert space of real random variables with finite second moments, defined on the probability space  $(\Omega, F, P)$ . In the following, we are interested in testing Granger causality in expectiles between  $Y$  and  $Z$ .

We now need to define the information sets as this is important for causality analysis. Hereafter, we consider a sequence  $I$  of “reference information sets” such that:

$$I = \{I(t) : t \in \mathbb{Z}, t > \omega\}, \text{ with } t < t' \Rightarrow I(t) \subseteq I(t'), \text{ for all } t > \omega,$$

where  $I(t)$  is an information set that represent a Hilbert subspace of  $L^2$ ,  $\omega \in \mathbb{Z} \cup \{-\infty\}$  represents a “starting point”, and  $\mathbb{Z}$  is the set of integers. The “starting point”  $\omega$  is typically equal to a finite initial date (such as  $\omega = -1, 0$  or  $1$ ) or to  $-\infty$ ; in the latter case  $I(t)$  is defined for all  $t \in \mathbb{Z}$ . The information set  $I(t)$  could correspond to a (possibly empty) set, whose elements represent the information available at any point of time, such as time independent variables (e.g., the constant in a regression model) and deterministic processes (e.g., deterministic trends).

We denote  $Y(\omega, t - 1]$  the information set spanned by  $Y_s$  for  $\omega < s \leq t - 1$ , and similarly for  $Z(\omega, t - 1]$ . That is,  $Y(\omega, t - 1]$  and  $Z(\omega, t - 1]$  represent the information contained in the history of the variables  $Y$  and  $Z$  up to time  $t - 1$ , respectively. Furthermore, the information sets obtained by “adding”  $Y(\omega, t - 1]$  to  $I(t - 1)$  and  $Z(\omega, t - 1]$  to  $I_Y(t - 1)$  are defined as

$$I_Y(t - 1) = I(t - 1) + Y(\omega, t - 1], \quad I_{YZ}(t - 1) = I_Y(t - 1) + Z(\omega, t - 1].$$

We assume that  $Y$  and  $Z$  are Markovians of orders  $p$  and  $q$ , respectively. In this case,  $Y(\omega, t - 1] \equiv Y(t - p, t - 1] \in \mathbb{R}^p$  and  $Z(\omega, t - 1] \equiv Z(t - q, t - 1] \in \mathbb{R}^q$ , and  $I_{YZ}(t - 1) \in \mathbb{R}^d$ , for  $d = p + q$ . Furthermore, let  $F_Y(y_t | I_{YZ}(t - 1))$  and  $F_Y(y_t | I_Y(t - 1))$  be the conditional distribution functions of  $y_t$  given  $I_{YZ}(t - 1)$  and  $I_Y(t - 1)$ , respectively, which we assume continuous for all  $y_t \in \mathbb{R}$ .

Before defining Granger causality in expectiles, we remind the reader how Granger non-causality in distribution and mean can be characterized in terms of restricted and unrestricted information sets. To avoid unnecessary repetitions, we treat only the causality from  $Z$  to  $Y$ . Following Granger (1969), causality from  $Z$  to  $Y$  one period ahead is defined as follows:  $Z$  causes  $Y$  if observations on  $Z$  up to time  $t - 1$  can help predict  $Y_t$  given the past of  $Y$  up to time  $t - 1$ . Formally, Granger non-causality from  $Z$  to  $Y$  is characterized as follows:

$$F_Y(y_t | I_{YZ}(t - 1)) = F_Y(y_t | I_Y(t - 1)), \quad \text{for all } y_t \in \mathbb{R}, \quad (1)$$

and will be referred to as Granger non-causality in *distribution*. Although the concept of Granger causality is naturally defined in terms of conditional distributions, applied work usually test for Granger non-causality in mean, which is only a necessary condition of Granger non-causality in distribution. In other words, the most used characterisation of Granger non-causality from  $Z$  to  $Y$  uses expectations:

$$E(y_t | I_{YZ}(t - 1)) = E(y_t | I_Y(t - 1)), \quad (2)$$

where  $E(y_t|I_{YZ}(t-1))$  and  $E(y_t|I_Y(t-1))$  denote the expectations of  $F_Y(y_t|I_{YZ}(t-1))$  and  $F_Y(y_t|I_Y(t-1))$ , respectively. Conditional mean, however, only measures one aspect of the conditional distribution and helps detect the mean dependence only. Indeed, a causal effect that concerns the tail area of the distribution might significantly differ from a causal effect that take place at the center of the distribution.

On the one hand, failing to reject non-causality in (2) is compatible with Granger non-causality in mean, but says nothing about Granger causality in other moments or other aspects (quantiles/expectiles) of conditional distribution. On the other hand, testing Granger non-causality in distribution using (1) will not be informative about the level(s) (mean, variance, other high-order moments, quantiles/expectiles) of distribution where the causality exists. To overcome this issue, we propose to use expectiles as a different way of looking at Granger causality between random variables. Expectiles will allow us to determine the pattern of causality across the conditional distribution of the variable of interest. They also provide sufficient condition for testing Granger non-causality in distribution as in (1), since expectiles are known to completely characterize the distribution. Formally, let  $\mu_\tau(.|I_{YZ}(t-1))$  and  $\mu_\tau(.|I_Y(t-1))$  denote the  $\tau$ th-expectiles of  $F_Y(.|I_{YZ}(t-1))$  and  $F_Y(.|I_Y(t-1))$ , respectively. Thus, an alternative way for testing Granger non-causality in distribution in (1) is by testing

$$\mu_\tau(y_t|I_{YZ}(t-1)) = \mu_\tau(y_t|I_Y(t-1)), \text{ for all } \tau \in \mathcal{T} \subset [0, 1]. \quad (3)$$

where  $\mathcal{T}$  is a subset or of a full set of  $[0, 1]$ . If (3) holds, then we say that  $Z$  does not Granger cause  $Y$  in all expectiles, and consequently there is no causality in distribution from  $Z$  to  $Y$ . In the next section, we will also be testing (3) at one given  $\tau$  only, which corresponds to testing Granger non-causality at one given expectile. The need for working with expectiles is motivated by the limits of quantiles. Despite their strong intuitive appeal, quantiles are not always satisfactory. They can be criticized for being somewhat difficult to work with and compute as the corresponding loss function is not continuously differentiable, although modern efficient linear programming algorithms are available to help. Most importantly, quantiles are relatively inefficient against long-tailed distributions as they are based on absolute rather than squared loss minimization. However, as we will see in the next sections, it's much easier to work with expectiles both theoretically and computationally; see Efron (1991) and Sobotka et al. (2013), Schnabel and Eilers (2009); Sobotka and Kneib (2012) among others.

### 3 Testing non-causality in expectiles

In this section, we use expectile regressions to build statistical procedures for testing Granger non-causality in expectiles. To this end, we first consider parametric regression models to estimate the  $\tau$ th-expectile of  $F_Y(\cdot|I_t)$ , for  $I_t$  some information set and  $\tau \in (0, 1)$ . We assume that the conditional function of the  $\tau$ th-expectile, say  $\mu_\tau(y_t|\cdot)$ , is correctly specified and belongs to a family  $M$  of linear and parametric functions:

$$M = \{m(\cdot, \theta(\tau)) \mid \theta(\cdot) : \tau \longrightarrow \theta(\tau) \in \Theta \subset \mathbb{R}^{d+1}, \text{ for } \tau \in \mathcal{T} \subset [0, 1]\},$$

where  $\theta(\tau)$  is a vector of parameters that characterise the  $\tau$ th-expectile of  $Y$ .

Let  $y_{t-1,p} = [y_{t-1}, \dots, y_{t-p}]'$ ,  $z_{t-1,q} = [z_{t-1}, \dots, z_{t-q}]'$ ,  $x_{t-1} = [1, y_{t-1,p}, z_{t-1,q}]$ , and consider the following correctly specified  $\tau$ th-expectile regression:

$$\begin{aligned} y_t &= \mu_\tau(y_t|I_{YZ}(t-1)) + \epsilon_t^\tau = a(\tau) + y'_{t-1,p}\alpha(\tau) + z'_{t-1,q}\beta(\tau) + \epsilon_t^\tau \\ &= x'_{t-1}\theta(\tau) + \epsilon_t^\tau, \quad \text{for } \tau \in (0, 1), \end{aligned} \quad (4)$$

where  $\theta(\tau) = [a(\tau), \alpha(\tau)', \beta(\tau)']'$  is the  $(d+1)$ -dimensional parameter vector, with  $d = p + q$ . The conditional  $\tau$ th-expectile of the error term  $\epsilon_t^\tau$  is equal zero as a result of correctly specified regression model. Using the parameters of regression (4), testing Granger non-causality from  $Z$  to a given  $\tau$ th-expectile of  $Y$  is equivalent to testing the null hypothesis:

$$H_0^\tau : \beta(\tau) = 0, \text{ for a given } \tau \in (0, 1), \quad (5)$$

and if the null is rejected this would suggest that  $Z$  Granger causes  $Y$  at a given expectile  $\tau$ .

In order to implement the test of causality in expectile, the vector of parameters  $\theta(\tau)$  is estimated by minimizing the following asymmetrically weighted absolute deviation function:

$$\hat{\theta}(\tau) = \underset{\theta(\tau) \in \mathbb{R}^{d+1}}{\operatorname{argmin}} \sum_{t=1}^T \psi_\tau(\epsilon_t^\tau) (y_t - x'_{t-1}\theta(\tau))^2, \quad (6)$$

where  $\psi_\tau(\epsilon_t^\tau) = |\tau - \mathbb{I}(\epsilon_t^\tau \leq 0)|$ , with  $\mathbb{I}(\cdot)$  is an indicator function that takes the value one if  $\epsilon_t^\tau \leq 0$  and zero otherwise. The minimization problem in (6) is known as expectile regression and the estimator  $\hat{\theta}(\tau)$  can also be obtained using a likelihood-based approach that uses Gaussian density with unequal weights placed on positive and negative disturbances. One of the advantages of expectile regression is that estimation basically reduces to (iteratively) weighted least

squares fits since the optimality criterion is differentiable, while linear programming routines have to be used in the case of quantile regression. Sobotka et al. (2011) compare expectiles and quantiles using both theory and simulations and find that the estimation of expectiles may be more efficient than the estimation of quantiles for a number of distributions. They also show a smaller probability to obtain crossing expectile curves than crossing quantile curves. In addition, contrary to quantile regression estimator, expectile regression estimator has an explicit form, which is given by:

$$\hat{\theta}(\tau) = \left( \sum_{t=1}^T \psi_{\tau}(\hat{\epsilon}_t^{\tau}) x_{t-1} x'_{t-1} \right)^{-1} \left( \sum_{t=1}^T \psi_{\tau}(\hat{\epsilon}_t^{\tau}) x_{t-1} y_t \right), \quad (7)$$

where  $\hat{\epsilon}_t^{\tau} = y_t - x'_{t-1} \hat{\theta}(\tau)$ .

Newey and Powell (1987) have studied the properties of the estimation of expectile regression

$$y_t = x'_{t-1} \theta(\tau) + \epsilon_t^{\tau}$$

when the data  $\{(x_{t-1}, y_t) \in \mathbb{R}^{d+1} \times \mathbb{R}\}$  is i.i.d. In particular, they showed that the least asymmetrically weighted squares estimate  $\hat{\theta}(\tau)$  is consistent and asymptotically normally distributed

$$\hat{\theta}(\tau) \sim N(\theta(\tau), \Sigma^{-1} V \Sigma^{-1})$$

with

$$\Sigma = \mathbb{E} [\psi_{\tau}(\epsilon_t^{\tau}) x_{t-1} x'_{t-1}] \quad \text{and} \quad V = \mathbb{E} [\psi_{\tau}^2(\epsilon_t^{\tau}) (\epsilon_t^{\tau})^2 x_{t-1} x'_{t-1}].$$

To estimate the variance-covariance matrix of  $\hat{\theta}(\tau)$ , Newey and Powell (1987) proposed to estimate  $\Sigma$  and  $V$  as follows:

$$\hat{\Sigma} = \frac{1}{T} \sum_{t=1}^T \psi_{\tau}(\hat{\epsilon}_t^{\tau}) x_{t-1} x'_{t-1}, \quad \hat{V} = \frac{1}{T} \sum_{t=1}^T x_{t-1} x'_{t-1} \psi_{\tau}(\hat{\epsilon}_t^{\tau})^2 \text{Var}(\hat{\epsilon}_t^{\tau}),$$

where  $\hat{\epsilon}_t^{\tau} = y_t - x'_{t-1} \hat{\theta}(\tau)$ . They then extended this result to establish the asymptotic normality of an estimator  $(\hat{\theta}(\tau_1), \dots, \hat{\theta}(\tau_m))$  for a grid point  $(\tau_1, \dots, \tau_m)$ , which was in turn used to build a test statistic for testing null hypotheses of the form  $R\theta(\tau) = \beta(\tau)$ , where  $R$  a selection matrix.

However, the data in our expectile regression model (4) is not i.i.d, thus testing the null in (5) or more generally

$$H_0 : \beta(\tau) = 0, \quad \forall \tau \in (0, 1) \quad (8)$$

requires weak convergence of the process  $\sqrt{T}(\hat{\theta}(\tau) - \theta(\tau))$ . To overcome this issue, we establish the weak convergence of the process  $\sqrt{T}(\hat{\theta}(\tau) - \theta(\tau))$  under time series data and we derive the asymptotic distribution of a test statistic that we use to test (8). The following assumptions are required to establish the asymptotic results:

**Assumptions:**

A.1  $\{(Y_t, Z_t), t \geq 0\}$  is a strictly stationary process.

A.2  $\epsilon_t^\tau$  are random variables with mean zero and variance  $\sigma^2$ .

A.3  $E[\psi_{\tau_1}(\epsilon_t^{\tau_1})\psi_{\tau_2}(\epsilon_t^{\tau_2})\epsilon_t^{\tau_1}\epsilon_t^{\tau_2}x_{t-1}x'_{t-1}]$  is nonsingular and  $E[\psi_{\tau}(\epsilon_t^\tau)x_{t-1}x'_{t-1}]$  is finite.

All of the above assumptions are fairly common in the literature. Assumption **A.2** on homoscedasticity of the error term  $\epsilon_t^\tau$  will be relaxed in the Monte Carlo simulation study. The following Theorem 1 establishes the asymptotic properties of the process  $\sqrt{T}(\hat{\theta}(\tau) - \theta(\tau))$  when the above assumptions are satisfied [see the proof of Theorem 1 in the appendix].

**Theorem 1** *Under Assumptions A.1-A.3, the empirical process  $\sqrt{T}(\hat{\theta}(\tau) - \theta(\tau))$  converges to a Gaussian process  $\mathcal{Z}(\tau)$  with covariance function*

$$Cov(\mathcal{Z}(\tau_1), \mathcal{Z}(\tau_2)) = \Psi_{\tau_1}^{-1}\Psi_{\tau_1, \tau_2}\Psi_{\tau_2}^{-1}, \quad (9)$$

where

$$\Psi_{\tau_i} = E[\psi_{\tau_i}(\epsilon_t^{\tau_i})x_{t-1}x'_{t-1}], \text{ for } i = 1, 2, \text{ and } \Psi_{\tau_1, \tau_2} = E[\psi_{\tau_1}(\epsilon_t^{\tau_1})\psi_{\tau_2}(\epsilon_t^{\tau_2})\epsilon_t^{\tau_1}\epsilon_t^{\tau_2}x_{t-1}x'_{t-1}].$$

Theorem 1 will be needed to establish the asymptotic distribution of a Kolmogorov-Smirnov type-Wald test statistic that we are going to use when we test (8). However, notice that the terms  $\Psi_{\tau_i}$ , for  $i = 1, 2$ , and  $\Psi_{\tau_1, \tau_2}$  in (9) are unknown, but they can be estimated as follows:

$$\hat{\Psi}_{\tau_i} = \frac{1}{T} \sum_{t=1}^T \psi_{\tau_i}(\hat{\epsilon}_t^{\tau_i})x_{t-1}x'_{t-1}, \text{ for } i = 1, 2, \text{ and } \hat{\Psi}_{\tau_1, \tau_2} = \frac{1}{T} \sum_{t=1}^T x_{t-1}x'_{t-1}\psi_{\tau_1}(\hat{\epsilon}_t^{\tau_1})\psi_{\tau_2}(\hat{\epsilon}_t^{\tau_2})\hat{\epsilon}_t^{\tau_1}\hat{\epsilon}_t^{\tau_2}. \quad (10)$$

For i.i.d. data, Newey and Powell (1987) state that  $\hat{\Psi}_{\tau_1}^{-1}\hat{\Psi}_{\tau_1, \tau_2}\hat{\Psi}_{\tau_2}^{-1}$  converges in probability to  $\Psi_{\tau_1}^{-1}\Psi_{\tau_1, \tau_2}\Psi_{\tau_2}^{-1}$ . The proof of this convergence can be adapted to time series under Assumptions **A.1-A.3**.

Now, from Theorem 1 and by observing that  $\sqrt{T}(\hat{\beta}(\tau) - \beta(\tau)) = \sqrt{T}(R\hat{\theta}(\tau) - R\theta(\tau))$ , for  $R = [0_{q,1+p}, I_{q,q}]$  a  $q \times (d+1)$  matrix, we can deduce that the process  $\sqrt{T}(\hat{\beta}(\tau) - \beta(\tau))$  converges to a Gaussian process  $\mathcal{W}(\tau)$  with covariance matrix:

$$Cov(\mathcal{W}(\tau_1), \mathcal{W}(\tau_2)) = \Omega_{\tau_1}^{-1} \Omega_{\tau_1, \tau_2} \Omega_{\tau_2}^{-1},$$

where

$$\Omega_{\tau_i} = E[\psi_{\tau_i}(\epsilon_t^{\tau_i}) z_{t-1,q} z'_{t-1,q}], \text{ for } i = 1, 2, \text{ and } \Omega_{\tau_1, \tau_2} = E[\psi_{\tau_1}(\epsilon_t^{\tau_1}) \psi_{\tau_2}(\epsilon_t^{\tau_2}) \epsilon_t^{\tau_1} \epsilon_t^{\tau_2} z_{t-1,q} z'_{t-1,q}].$$

By analogy to (10),  $\Omega_{\tau_i}$  and  $\Omega_{\tau_1, \tau_2}$  can be estimated as:

$$\hat{\Omega}_{\tau_i} = \frac{1}{T} \sum_{t=1}^T \psi_{\tau}(\hat{\epsilon}_t^{\tau_i}) z_{t-1,q} z'_{t-1,q}, \text{ for } i = 1, 2, \text{ and } \hat{\Omega}_{\tau_1, \tau_2} = \frac{1}{T} \sum_{t=1}^T z_{t-1,q} z'_{t-1,q} \psi_{\tau_1}(\hat{\epsilon}_t^{\tau_1}) \psi_{\tau_2}(\hat{\epsilon}_t^{\tau_2}) \hat{\epsilon}_t^{\tau_1} \hat{\epsilon}_t^{\tau_2}. \quad (11)$$

In Corollary 1 below, we derive the asymptotic normality of  $\sqrt{T}(\hat{\beta}(\tau) - \beta(\tau))$  for fixed  $\tau \in (0, 1)$ . The proof of the Corollary 1 can be straightforwardly deduced from Theorem 1, hence it will be omitted. Moreover, the result in this corollary will be used to build a test for

$$H_0^{\tau} : \beta(\tau) = 0 \quad \text{versus} \quad H_1^{\tau} : \beta(\tau) \neq 0.$$

**Corollary 1.** *For a fixed  $\tau$ , for  $\tau \in (0, 1)$ , and under Assumptions **A.1-A.3**, we have*

$$\sqrt{T} \left( \hat{\Omega}_{\tau}^{-1} \hat{\Omega}_{\tau, \tau} \hat{\Omega}_{\tau}^{-1} \right)^{-1/2} (\hat{\beta}(\tau) - \beta(\tau))$$

*converges to  $N(0, I_{q \times q})$ , where  $\hat{\Omega}_{\tau}$  and  $\hat{\Omega}_{\tau, \tau}$  are similarly defined in (11).*

To test the null hypothesis in (8), we first define the empirical process:

$$W_T(\tau) = T(\hat{\beta}(\tau) - \beta(\tau))' \left( \hat{\Omega}_{\tau}^{-1} \hat{\Omega}_{\tau, \tau} \hat{\Omega}_{\tau}^{-1} \right)^{-1} (\hat{\beta}(\tau) - \beta(\tau)).$$

Using the process  $W_T(\tau)$ , we consider a test statistic that is based on Kolmogorov-Smirnov type-Wald test, say  $KSW_T$ , which is defined as follows:

$$KSW_T = \sup_{\tau \in (0,1)} W_T(\tau).$$

The following proposition states the asymptotic distribution of the test  $KSW_T$  for testing (8). The proof of Proposition 1 will be omitted as it can be straightforwardly deduced from Corollary 1.

**Proposition 1** *Under Assumptions A.1-A.3, and the null hypothesis  $H_0$ , we have*

$$KSW_T \xrightarrow{d} \sup_{\tau} \|\mathcal{Z}(\tau)\|_2^2,$$

where  $\mathcal{Z}$  is a Gaussian process and  $\|\cdot\|_2$  is the Euclidean norm. For fixed  $\tau$ ,  $\|\mathcal{Z}(\tau)\|_2^2$  is a centred Chi-square random variable with  $q$  degree of freedom.

The result in Proposition 1 will be used to test Granger causality in distribution from  $Z$  to  $Y$ . We next use Monte Carlo simulations to assess the finite-sample properties (size and power) of the tests derived in Corollary 1 and Proposition 1 for a variety of data-generating processes and different sample sizes. We will also compare the properties of our expectiles-based test with the existing quantiles-based test, see e.g. Koenker and Machado (1999).

## 4 Monte Carlo simulations

We conduct a Monte Carlo simulation study to investigate the performance of the tests we proposed previously. Our primary interest is to assess the properties of the tests derived in Corollary 1 and Proposition 1. In particular, we examine their size and power properties using the data-generating processes (DGPs) presented in Table 1.

The first four DGPs [DGP1-DGP4] in Table 1 were used to evaluate the empirical size of the tests since the causality in these DGPs does not exist. DGP3 and DGP4 allow for heteroscedasticity, with the former model includes the ARCH effect of Engle (1982) and the latter incorporates the GARCH effect of Bollerslev (1986). The last three DGPs [DGP5-DGP7] of Table 1 allow for Granger causality in expectiles, thus they serve to illustrate the power of the tests. DGP5 exhibits Granger causality in the presence of homoscedastic errors, whereas DGP6 and DGP7 both allow for heteroskedastic errors.

These DGPs focus on testing Granger causality from  $Z$  to  $Y$  using the regression equation:

$$Y_t = \mu + 0.5Y_{t-1} + c(\tau)Z_{t-1} + \epsilon_{1t},$$

where  $Z$  is generated using the processes in Table 1. We then test the null hypothesis  $H_0: c(\tau) = 0$  against the alternative hypothesis  $H_1: c(\tau) \neq 0$ , for  $\tau \in (0, 1)$ . In our simulation and under the alternative hypothesis,  $c(\tau)$  does not change with  $\tau$  and takes one of the following

Table 1: DGPs used in the simulation study

DGPs	$Y_t$	$Z_t$
DGP1	$\epsilon_{1t}$	$\epsilon_{2t}$
DGP2	$Y_t = 0.5Y_{t-1} + \epsilon_{1t}$	$Z_t = 0.5Z_{t-1} + \epsilon_{2t}$
DGP3	$Y_t = (0.01 + 0.5Y_{t-1}^2)^{0.5}\epsilon_{1t}$	$Z_t = 0.5Z_{t-1} + \epsilon_{2t}$
DGP4	$Y_t = \sqrt{h_{1,T}(\delta)}\epsilon_{1t}$ , with $h_{1,T}(\delta) = 0.01 + 0.9h_{1,t-1} + 0.05Y_{t-1}^2$	$Z_t = \sqrt{h_{2,T}(\delta)}\epsilon_{2t}$ , with $h_{2,T}(\delta) = 0.01 + 0.9h_{2,t-1} + 0.05Z_{t-1}^2$
DGP5	$Y_t = 0.5Y_{t-1} + cZ_{t-1} + \epsilon_{1t}$	$Z_t = 0.5Z_{t-1} + \epsilon_{2t}$
DGP6	$Y_t = 0.5Y_{t-1} + cZ_{t-1} + \epsilon_{1t}$ $\epsilon_{1t} \sim N(0, t/T + 2)$	$Z_t = 0.5Z_{t-1} + \epsilon_{2t}$ $\epsilon_{2t} \sim N(0, t/T + 2)$
DGP7	$Y_t = 0.5Y_{t-1} + cZ_{t-1} + \epsilon_{1t}$ $\epsilon_{1t} \sim N(0, 1/t + 2)$	$Z_t = 0.5Z_{t-1} + \epsilon_{2t}$ $\epsilon_{2t} \sim N(0, 1/t + 2)$

**Note:** This table summarizes the DGPs that we consider in the simulation study to investigate the properties (size and power) of the tests of Granger causality in expectiles.

values  $c(\tau) = 0.1, 0.5,$  and  $0.9,$  which capture different degrees of causality from  $Z$  to  $Y$ . A higher (in absolute value) of  $c(\tau)$  indicates a stronger causality from  $Z$  to  $Y$ .

Four sample sizes  $T = 100, 200, 500,$  and  $1000$  were considered. For each sample size  $T$  and DGP, 1000 (number of simulations) independent realisations of length  $T$  were obtained as follows: (i) we generate  $T + 200$  i.i.d. noises  $\epsilon_{1t}$  and  $\epsilon_{2t}$  from  $N(0, 1),$  except for DGP6 and DGP7 that are generated under normality with variances different from one, and (ii) each noise's sequence is used to generate  $Y_t$  and  $Z_t,$  for  $t = 1, \dots, T + 200.$  The initial values of  $Y_t$  and  $Z_t$  were set to zero (resp. to one). To attenuate the impact of these initial values, the first 200 observations were discarded.

#### 4.1 Size and power of expectile-based tests

The results for the empirical size of the test derived in Corollary 1 are presented in Table 2. From this, we see that for almost all DGPs and sample sizes under consideration, the empirical sizes of the test of Granger causality at a given expectile are very close to the nominal levels  $\alpha = 5\%$  and  $\alpha = 10\%.$  The empirical size is well controlled in both small and large samples and in the presence and absence of heteroskedasticity. This result generally holds for extreme expectiles like for  $\tau = 0.1$  and  $0.9.$

The results for the empirical power of the test in Corollary 1 are presented in Tables 3-5. The latter show that the proposed test has good power (very close to 1 or equal to 1) for all expectiles  $\tau,$  for  $\tau \in (0, 1),$  all DGPs and sample sizes we considered, particularly when the degree of causality is not low (e.g.  $c = 0.5$  and  $c = 0.9).$  When the degree of causality is low, say  $c = 0.1,$  then as expected the power is low, but it improves significantly (above 0.9) once the sample size is increased to say  $T = 1000.$

Finally, Tables 6 and 7 provide the results for the empirical size and power of the Kolmogorov-Smirnov type-Wald test in Proposition 1. The results show that the latter controls well its size and has a very good power for a variety of data-generating processes and different sample sizes. The power reaches 1 even for small samples, e.g.  $T = 100.$

Table 2: Empirical size of the proposed test of the Granger non-causality test in expectiles

DGPs	$\alpha = 5\%$				$\alpha = 10\%$			
	DGP1	DGP2	DGP3	DGP4	DGP1	DGP2	DGP3	DGP4
$T = 100$								
$\tau = 0.1$	0.043	0.075	0.058	0.062	0.10	0.11	0.11	0.117
$\tau = 0.25$	0.054	0.061	0.053	0.052	0.11	0.114	0.109	0.107
$\tau = 0.5$	0.045	0.064	0.056	0.046	0.10	0.106	0.091	0.10
$\tau = 0.75$	0.059	0.065	0.067	0.053	0.09	0.099	0.112	0.115
$\tau = 0.9$	0.054	0.053	0.071	0.054	0.9	0.103	0.101	0.111
$T = 200$								
$\tau = 0.1$	0.058	0.061	0.058	0.053	0.112	0.097	0.112	0.114
$\tau = 0.25$	0.046	0.065	0.045	0.059	0.095	0.097	0.106	0.108
$\tau = 0.5$	0.053	0.052	0.051	0.049	0.112	0.119	0.091	0.104
$\tau = 0.75$	0.055	0.059	0.065	0.048	0.111	0.094	0.100	0.101
$\tau = 0.9$	0.051	0.066	0.048	0.049	0.114	0.093	0.102	0.092
$T = 500$								
$\tau = 0.1$	0.047	0.051	0.068	0.049	0.101	0.103	0.120	0.116
$\tau = 0.25$	0.051	0.054	0.055	0.056	0.091	0.095	0.104	0.115
$\tau = 0.5$	0.054	0.048	0.057	0.061	0.104	0.096	0.09	0.104
$\tau = 0.75$	0.055	0.052	0.064	0.053	0.104	0.082	0.10	0.104
$\tau = 0.9$	0.054	0.046	0.057	0.056	0.083	0.106	0.110	0.101
$T = 1000$								
$\tau = 0.1$	0.055	0.05	0.057	0.06	0.113	0.097	0.110	0.095
$\tau = 0.25$	0.057	0.06	0.061	0.04	0.099	0.110	0.113	0.103
$\tau = 0.5$	0.051	0.057	0.048	0.052	0.104	0.089	0.102	0.105
$\tau = 0.75$	0.051	0.045	0.052	0.043	0.093	0.105	0.101	0.09
$\tau = 0.9$	0.045	0.056	0.065	0.046	0.102	0.120	0.110	0.120

**Note:** This table reports the empirical size of the test in Corollary 1 for testing the Granger non-causality at a given expectile from  $Z$  to  $Y$  at  $\alpha = 5\%$  and  $10\%$  significance levels. The number of simulations is equal to 1000.

Table 3: Empirical power of the proposed test of the Granger non-causality test in expectiles

<b>DGP5</b>								
Sample size	$T = 100$		$T = 200$		$T = 500$		$T = 1000$	
	$\alpha = 5\%$	$\alpha = 10\%$						
<b>c=0.1</b>								
$\tau = 0.10$	0.178	0.229	0.253	0.375	0.538	0.692	0.833	0.903
$\tau = 0.25$	0.194	0.263	0.31	0.448	0.670	0.766	0.927	0.962
$\tau = 0.50$	0.211	0.327	0.361	0.474	0.713	0.818	0.952	0.981
$\tau = 0.75$	0.202	0.290	0.345	0.440	0.695	0.775	0.938	0.961
$\tau = 0.90$	0.167	0.251	0.238	0.364	0.522	0.673	0.835	0.907
<b>c=0.5</b>								
$\tau = 0.10$	0.975	0.996	1	1	1	1	1	1
$\tau = 0.25$	0.997	1	1	1	1	1	1	1
$\tau = 0.50$	1	1	1	1	1	1	1	1
$\tau = 0.75$	0.999	0.998	1	1	1	1	1	1
$\tau = 0.90$	0.977	0.987	1	1	1	1	1	1
<b>c=0.9</b>								
$\tau = 0.10$	1	1	1	1	1	1	1	1
$\tau = 0.25$	1	1	1	1	1	1	1	1
$\tau = 0.50$	1	1	1	1	1	1	1	1
$\tau = 0.75$	1	1	1	1	1	1	1	1
$\tau = 0.90$	1	1	1	1	1	1	1	1

**Note:** This table reports the empirical power of the test in Corollary 1 for testing the Granger non-causality at a given expectile from  $Z$  to  $Y$  at  $\alpha = 5\%$  and  $10\%$  significance levels. The number of simulations is equal to 1000.

Table 4: Empirical power of the proposed test of the Granger non-causality test in expectiles

<b>DGP6</b>								
Sample size	$T = 100$		$T = 200$		$T = 500$		$T = 1000$	
	$\alpha = 5\%$	$\alpha = 10\%$						
<b>c=0.1</b>								
$\tau = 0.10$	0.152	0.260	0.277	0.345	0.539	0.661	0.829	0.883
$\tau = 0.25$	0.203	0.306	0.303	0.448	0.663	0.763	0.920	0.948
$\tau = 0.50$	0.201	0.284	0.354	0.478	0.691	0.802	0.952	0.963
$\tau = 0.75$	0.188	0.292	0.307	0.445	0.672	0.760	0.920	0.962
$\tau = 0.90$	0.147	0.257	0.253	0.341	0.529	0.634	0.838	0.885
<b>c=0.5</b>								
$\tau = 0.10$	0.968	0.989	0.998	1	1	1	1	1
$\tau = 0.25$	0.998	0.998	1	1	1	1	1	1
$\tau = 0.50$	0.999	1	1	1	1	1	1	1
$\tau = 0.75$	0.997	0.998	1	1	1	1	1	1
$\tau = 0.90$	0.965	0.991	1	1	1	1	1	1
<b>c=0.9</b>								
$\tau = 0.10$	1	1	1	1	1	1	1	1
$\tau = 0.25$	1	1	1	1	1	1	1	1
$\tau = 0.50$	1	1	1	1	1	1	1	1
$\tau = 0.75$	1	1	1	1	1	1	1	1
$\tau = 0.90$	1	1	1	1	1	1	1	1

**Note:** This table reports the empirical power of the test in Corollary 1 for testing the Granger non-causality at a given expectile from  $Z$  to  $Y$  at  $\alpha = 5\%$  and  $10\%$  significance levels. The number of simulations is equal to 1000.

Table 5: Empirical power of the proposed test of the Granger non-causality test in expectiles

<b>DGP7</b>								
Sample size	$T = 100$		$T = 200$		$T = 500$		$T = 1000$	
	$\alpha = 5\%$	$\alpha = 10\%$						
<b>c=0.1</b>								
$\tau = 0.10$	0.161	0.236	0.262	0.386	0.556	0.644	0.834	0.903
$\tau = 0.25$	0.176	0.288	0.337	0.423	0.660	0.781	0.927	0.963
$\tau = 0.50$	0.215	0.310	0.350	0.485	0.726	0.789	0.950	0.984
$\tau = 0.75$	0.215	0.309	0.326	0.437	0.693	0.768	0.941	0.970
$\tau = 0.90$	0.174	0.239	0.222	0.361	0.533	0.687	0.839	0.904
<b>c=0.5</b>								
$\tau = 0.10$	0.976	0.991	1	1	1	1	1	1
$\tau = 0.25$	0.996	1	1	1	1	1	1	1
$\tau = 0.50$	1	1	1	1	1	1	1	1
$\tau = 0.75$	1	1	1	1	1	1	1	1
$\tau = 0.90$	0.978	0.991	1	1	1	1	1	1
<b>c=0.9</b>								
$\tau = 0.10$	1	1	1	1	1	1	1	1
$\tau = 0.25$	1	1	1	1	1	1	1	1
$\tau = 0.50$	1	1	1	1	1	1	1	1
$\tau = 0.75$	1	1	1	1	1	1	1	1
$\tau = 0.90$	1	1	1	1	1	1	1	1

**Note:** This table reports the empirical power of the test in Corollary 1 for testing the Granger non-causality at a given expectile from  $Z$  to  $Y$  at  $\alpha = 5\%$  and  $10\%$  significance levels. The number of simulations is equal to 1000.

Table 6: Empirical size of the proposed sup-Wald test of Granger non-causality in expectiles


	$\alpha = 5\%$				$\alpha = 10\%$			
	<b>DGP1</b>	<b>DGP2</b>	<b>DGP3</b>	<b>DGP4</b>	<b>DGP1</b>	<b>DGP2</b>	<b>DGP3</b>	<b>DGP4</b>
$T = 100$	0.044	0.048	0.046	0.058	0.088	0.100	0.096	0.102
$T = 200$	0.048	0.046	0.054	0.064	0.082	0.088	0.114	0.110
$T = 500$	0.062	0.066	0.038	0.046	0.110	0.136	0.098	0.104
$T = 1000$	0.050	0.044	0.038	0.046	0.102	0.100	0.096	0.104

**Note:** This table reports the empirical size of the test in Proposition 1 for testing Granger non-causality in expectiles from  $Z$  to  $Y$  at  $\alpha = 5\%$  and  $10\%$  significance level. The number of simulations is equal to 1000.


Table 7: Empirical power of our sup-Wald test of Granger non-causality in expectiles

	$\alpha = 5\%$			$\alpha = 10\%$		
	<b>DGP5</b>	<b>DGP6</b>	<b>DGP7</b>	<b>DGP5</b>	<b>DGP6</b>	<b>DGP7</b>
$T = 100$	0.982	0.984	0.986	0.996	0.998	0.998
$T = 200$	1.000	1.000	1.000	1.000	1.000	1.000
$T = 500$	1.000	1.000	1.000	1.000	1.000	1.000
$T = 1000$	1.000	1.000	1.000	1.000	1.000	1.000


**Note:** This table reports the empirical power of the test in Proposition 1 for testing Granger non-causality in expectiles from  $Z$  to  $Y$  at  $\alpha = 5\%$  and  $10\%$  significance level. The number of simulations is equal to 1000.




(a)  $\tau = 75\%$



(b)  $\tau = 50\%$



(c)  $\tau = 25\%$



(d)  $\tau = 10\%$

**Figure 1:** This figure compares the power of expectile-based test (Corollary 1) with the power of quantile-based test [see e.g. Koenker and Machado (1999)] for testing Granger non-causality from  $Z$  to  $Y$  at  $\alpha = 5\%$  significance level and for a range of values of  $\tau$  :  $\tau = 75\%$ ,  $50\%$ ,  $25\%$ , and  $10\%$ . The number of simulations is equal to 1000.

## 4.2 Expectiles versus quantiles-based tests

We consider an additional simulation exercise to compare the size and power of the expectile-based test in Corollary 1 with those of quantile-based test [see e.g. Koenker and Machado (1999)] for different values of  $\tau$ . We use the same simulation settings as in the previous subsection and for a fair comparison between the two tests, we consider the DGP5 of Table 1:

$$Y_t = \mu + 0.5Y_{t-1} + cZ_{t-1} + \epsilon_{1t},$$

where  $Z_t = 0.5Z_{t-1} + \epsilon_{2t}$ , with  $\epsilon_{1t}, \epsilon_{2t} \sim N(0, 1)$ . Under DGP5, conditional expectiles and quantiles of  $Y_t$  can be expressed as follows:

$$\mu_\tau(Y_t|Y_{t-1}, Z_{t-1}) = \mu + 0.5Y_{t-1} + cZ_{t-1} + \mu_\tau(\epsilon_{1t}|Y_{t-1}, Z_{t-1}) \quad (12)$$

$$q_\tau(Y_t|Y_{t-1}, Z_{t-1}) = \mu + 0.5Y_{t-1} + cZ_{t-1} + q_\tau(\epsilon_{1t}|Y_{t-1}, Z_{t-1}),$$

for  $\tau \in (0, 1)$ , where  $\mu_\tau(\epsilon_{1t}|Y_{t-1}, Z_{t-1})$  and  $q_\tau(\epsilon_{1t}|Y_{t-1}, Z_{t-1})$  are the  $\tau$ th-expectile and  $\tau$ th-quantile of the error term  $\epsilon_{1t}$ . The two equations in (12) show that the degree of causality (measured by  $c$ ) is the same for expectiles and quantiles and does not change with  $\tau$ , which guarantees fair comparison of the properties of expectile- and quantile-based tests. The results are reported in Figures 1-4.

Figures 1-4 illustrate the powers of the above-mentioned tests for testing Granger non-causality based on expectile and quantile regressions for a range of values of  $\tau$ . From these, we see that the test derived in Corollary 1 outperforms the test based on quantile regression even when the degree of causality is weak ( $c$  low). All the figures show that our test not only outperforms the test based on quantiles for large samples like  $T = 500$ , but it also provides reliable results for smaller samples like  $T = 100$ . The results are consistent across all the considered values of  $\tau$ .

## 5 Empirical applications

### 5.1 Causality between returns and volume

As a first application of the tests proposed in Section (3), we consider the problem of testing Granger causality (hereafter predictability) in expectiles from volume to market returns. Predicting the conditional distribution of stock returns using mean and quantile regressions has been the focus of many empirical studies, see Fama and French (1988), Keim and Stambaugh (1986), Ülkü and Onishchenko (2019), Baur et al. (2012) among others. Chuang et al. (2009) have recently investigated the predictability of stock returns using volume based on parametric quantile regressions. They first define Granger non-causality in all quantiles and propose testing non-causality based on a sup-Wald type test. Using three major stock market indices, they find that the causal effects of volume on return are usually heterogeneous across quantiles. In particular, the quantile causal effects of volume on return exhibit a spectrum of V-shape

relations so that the dispersion of return distribution increases with lagged volume.

There is, however, no study on the predictability of stock return distribution using volume based on expectile regressions. Expectiles have the advantage of capturing similar information as quantiles, but they also have the merit of being much more straightforward to use than quantiles, since expectiles are defined as least squares analogue of quantiles. Furthermore, testing Granger causality (predictability) at all expectiles - as we do in this application and the next one - provides a sufficient condition for testing Granger causality in distribution. Thus, in this section we apply our tests to re-examine Granger causality in expectiles between returns and volume using daily data on three major stock market indices: NYSE, S&P 500 and FTSE 100.

### 5.1.1 Data description and results

The dataset that we use in this application comes from Yahoo Finance and consists of daily observations on NYSE, S&P 500 and FTSE 100 and their volumes. The sample runs from January 2010 to January 2020 for a total of 3032 observations. We compute the continuously compounded changes in prices (returns) and trading volume (volume growth rate) and we perform Augmented Dickey-Fuller tests (ADF-tests) for testing non-stationarity of the logarithmic price and volume and their first differences. Using ADF-tests with an intercept and trend, with an intercept and without trend, and without intercept and trend, the results show that all variables in logarithmic form are non-stationary, but their first differences are. Based on this outcome, we model the first difference of logarithmic price and volume rather than their levels. Consequently, the causality relations have to be interpreted in terms of growth rates.

To test the causality in expectiles from volume to stock market return, we use the expectile regression (ER):

$$r_t = \alpha_0(\tau) + \sum_{i=1}^q \alpha_i(\tau) r_{t-i} + \sum_{j=1}^q \beta_j(\tau) v_{t-j} + \epsilon_t^\tau,$$

where  $r_{t-j}$  is the continuously compounded return from one of the stock market indices (NYSE, S&P 500 and FTSE 100) at time  $t - j$ ,  $v_{t-j}$  is the corresponding continuously compounded volume at time  $t - j$ , and  $\epsilon_t^\tau$  is the error term with zero conditional expectile. Returns are calculated as  $r_t = 100 \times (\ln(p_t) - \ln(p_{t-1}))$  and the growth rate of the volume as  $v_t = 100 \times (\ln(V_t) - \ln(V_{t-1}))$ , where  $p_t$  and  $V_t$  are the price and traded share volume of a given index at

time  $t$ . We then test the null hypotheses:

$$H_0^\tau : \beta(\tau) = 0 \quad \text{versus} \quad H_1^\tau : \beta(\tau) \neq 0$$

and

$$H_0 : \beta(\tau) = 0, \forall \tau \in (0, 1) \quad \text{versus} \quad H_1 : \beta(\tau) \neq 0 \text{ for some or all } \tau,$$

where  $\beta(\tau) = (\beta_1(\tau), \dots, \beta_q(\tau))'$ . The former null  $H_0^\tau$  corresponds to testing non-causality (non-predictability) from volume to stock returns at a given expectile  $\tau$ , whereas the latter null  $H_0$  is about testing non-causality (non-predictability) at all expectiles  $\tau \in (0, 1)$ , which is equivalent to testing independence between volume and stock returns.

The results for testing  $H_0^\tau$  and  $H_0$  for different lags  $q$  are reported in Table 8. Three different values of the number of lags are considered:  $q = 3, 10, \text{ and } 12$ . Table 8 shows that changes in volume are causing changes in stock returns. The causality happens generally at the upper expectiles (0.9 to 1) of stock returns, except for FTSE. The Kolmogorov-Smirnov type-Wald test of Proposition 1 also confirms the causality from volume to stock returns of NYSE, S&P 500 and FTSE 100, especially when the number lags is high ( $q = 10, 12$ ).

## 5.2 Stock market return and exchange rates

Given its significance for the economy, the causal relationship between stock prices and exchange rates is of great importance for academics, policymakers and professionals. Thus, as a second application of our tests, we investigate the causality in expectiles from exchange rate to market returns.

In the literature, there is no academic consensus about this relationship and the results are somewhat mixed as to whether or not exchange rates affect the prices of stock indexes. On the one hand, Chow et al. (1997) find evidence that the link between changes in exchange rate and stock returns is stronger at long horizons, whereas Williamson (2001) find that this link has changed over time. Yang et al. (2014) have applied parametric regression models to investigate the causality between stock returns and exchange rate for nine Asian markets. Their empirical results show that there are more causal relations based on the quantile regression than the conventional mean regression. On the other hand, Griffin and Stulz (2001) conclude that this link is small and hardly significant. Most of the conclusions, however, were reached using mean

Table 8: Granger-causality in expectiles between stock market return and volume

$\tau$	Number of Lags = 3			Number of Lags = 10			Number of Lags = 12		
	NYSE	S&P	FTSE	NYSE	S&P	FTSE	NYSE	S&P	FTSE
0.01	1.132	3.865	3.606	9.716	13.01	<b>22.21**</b>	11.41	10.99	<b>21.28**</b>
0.05	0.921	1.401	2.564	6.313	6.287	7.820	8.141	8.37	9.070
0.1	0.874	1.475	2.745	6.882	7.036	7.613	8.552	8.31	10.27
0.2	1.187	1.934	3.277	5.841	7.991	10.82	7.809	8.72	11.55
0.3	1.826	2.342	3.191	6.114	8.004	12.75	8.169	10.05	13.36
0.4	2.401	2.693	3.004	6.637	7.913	14.08	8.641	10.04	14.59
0.5	3.056	3.163	2.703	7.069	8.303	14.86	9.424	10.54	15.46
0.6	3.890	3.776	2.251	7.648	9.070	15.00	10.445	11.58	15.56
0.7	4.768	4.354	1.800	8.520	10.07	15.21	11.76	13.02	15.86
0.8	5.523	5.334	1.445	10.29	12.05	15.27	15.31	15.15	17.02
0.9	<b>7.360*</b>	<b>6.410*</b>	1.638	14.62	15.96	14.12	<b>19.01*</b>	17.16	14.42
0.95	<b>8.196**</b>	<b>7.332*</b>	2.654	<b>21.58**</b>	<b>20.31**</b>	15.11	<b>25.94**</b>	18.24	15.14
0.999	<b>12.60***</b>	2.697	1.18	<b>51.74***</b>	<b>36.14***</b>	<b>19.18**</b>	<b>40.95***</b>	<b>30.04***</b>	15.46
Sup-Wald	<b>12.60***</b>	<b>7.332*</b>	3.606	<b>51.74***</b>	<b>36.14***</b>	<b>22.21**</b>	<b>40.95***</b>	<b>30.04***</b>	<b>21.28**</b>

**Note:** This table reports the results of Granger-causality in expectiles between stock market return (NYSE, S&P, FTSE) and volume. Each entry is a test for the null hypothesis of non-causality from volume to return at a given expectile  $\tau$ , for  $\tau \in (0, 1)$ . \*\*\*, \*\*, and \* denote significance at 1%, 5%, and 10% levels, respectively, and the corresponding critical values are 11.344, 7.814, and 6.251 for  $\chi^2(3)$  [Number of Lags = 3], and 23.209, 18.307, and 15.987 for  $\chi^2(10)$  [Number of Lags = 10], and 26.217, 21.026, and 18.549 for  $\chi^2(12)$  [Number of Lags = 12].

regression-based tests. In this section, we apply the Granger causality in expectiles tests to examine the causal relationship between stock prices and exchange rates.

### 5.2.1 Data description and results

For this second application, we use data on S&P 500 Index and US/Canada, US/UK and US/Japan exchange rates. The data sets consist of monthly observations on S&P 500 Index and exchange rates and their obtained from St. Louis Fed and Yahoo Finance, respectively, and the sample runs from January 2004 to January 2020 for a total of 193 observations. As in the first application, we perform ADF-tests for testing non-stationarity of the logarithmic price and exchange rates and their first differences. The results show that all variables in logarithmic form are non-stationary, whereas their first differences are stationary. Thus, we model the first differences of logarithmic price and exchange rates rather than their level.

To test the causality in expectiles from exchange rates to stock market return, we use the expectile regression:

$$r_t = \alpha_0(\tau) + \sum_{j=1}^q \alpha_j(\tau) r_{t-j} + \sum_{j=1}^q \beta_j(\tau) ex_{t-j}^{\text{US}/i} + \epsilon_t^\tau,$$

where  $ex_{t-j}^{\text{US}/i}$ , for  $i = \text{Canada, UK and Japan}$ , is the growth rate of exchange rate US/country  $i$  at time  $t-j$ ,  $r_{t-j}$  is the continuously compounded return from the S&P 500 index at time  $t-j$ , and  $\epsilon_t^\tau$  is the error term with zero conditional expectile. As before, returns are calculated as  $r_t = 100 \times (\ln(p_t) - \ln(p_{t-1}))$  and exchange rates as  $ex_{t-j}^{\text{US}/i} = 100 \times \left( \ln \left( EX_t^{\text{US}/i} \right) - \ln \left( EX_{t-1}^{\text{US}/i} \right) \right)$ , where  $p_t$  and  $EX_t^{\text{US}/i}$  are the price of S&P 500 index and exchange rate US/country  $i$  at time  $t$ , respectively. We then test the null hypotheses:

$$H_0^\tau : \beta(\tau) = 0 \quad \text{versus} \quad H_1^\tau : \beta(\tau) \neq 0$$

and

$$H_0 : \beta(\tau) = 0, \forall \tau \in (0, 1) \quad \text{versus} \quad H_1 : \beta(\tau) \neq 0 \text{ for some or all } \tau.$$

where  $\beta(\tau) = (\beta_1(\tau), \dots, \beta_q(\tau))'$ . The former null  $H_0^\tau$  corresponds to testing non-causality from growth rate of exchange rate to stock return at a given expectile  $\tau$ , whereas the latter null hypothesis  $H_0$  is about testing non-causality at all expectiles  $\tau \in (0, 1)$ , which is equivalent to testing independence between growth rate of exchange rate and stock return.

Table 9: Granger-causality in expectiles between stock market return and exchange rates

$\tau$	Number of Lags = 3			Number of Lags = 10			Number of Lags = 12		
	$EX_{CAD}$	$EX_{UK}$	$EX_{JAP}$	$EX_{CAD}$	$EX_{UK}$	$EX_{JAP}$	$EX_{CAD}$	$EX_{UK}$	$EX_{JAP}$
0.01	0.987	<b>8.271**</b>	<b>7.562*</b>	<b>30.67***</b>	5.091	13.16	16.64	<b>165.84***</b>	<b>22.59***</b>
0.05	0.610	<b>6.298*</b>	1.973	<b>26.73***</b>	12.28	10.74	<b>37.39***</b>	<b>63.03***</b>	11.10
0.1	0.913	<b>8.331**</b>	1.404	12.12	<b>21.61**</b>	8.635	<b>27.29***</b>	<b>49.02***</b>	10.47
0.2	1.143	<b>11.17**</b>	1.202	11.62	<b>21.42**</b>	8.293	<b>18.58*</b>	<b>43.47***</b>	9.524
0.3	1.127	<b>9.474**</b>	1.057	10.82	<b>18.49**</b>	8.642	16.52	<b>31.88***</b>	9.763
0.4	1.114	<b>8.207**</b>	0.927	11.12	<b>18.74**</b>	9.633	15.14	<b>26.30***</b>	9.594
0.5	1.076	<b>7.508*</b>	0.746	11.41	<b>16.72*</b>	8.495	14.39	<b>21.90*</b>	9.115
0.6	1.072	<b>6.861*</b>	0.581	11.95	14.96	8.130	13.53	<b>19.11*</b>	8.925
0.7	1.060	<b>6.387*</b>	0.371	12.89	14.11	7.836	13.90	17.91	8.923
0.8	0.94	6.008	0.216	14.23	12.48	8.147	14.78	16.03	8.861
0.9	0.547	4.232	0.106	17.83	11.05	7.656	13.22	12.15	7.734
0.95	0.571	4.262	0.012	11.11	16.13	7.502	14.67	14.87	6.497
0.99	1.631	5.328	2.308	<b>22.37**</b>	7.016	5.216	12.43	13.67	7.931
Sup-Wald	1.631	<b>11.33**</b>	<b>7.562*</b>	<b>30.67***</b>	<b>22.66**</b>	14.99	<b>57.31***</b>	<b>165.84***</b>	<b>22.59***</b>

**Note:** This table reports the results of Granger-causality in expectiles between stock market return and exchange rate ( $EX_{CAD}$ ,  $EX_{UK}$ ,  $EX_{JAP}$ ). Each entry is a test for the null hypothesis of non-causality from exchange rate to return at a given expectile  $\tau$ , for  $\tau \in (0, 1)$ . \*\*\*, \*\*, and \* denote significance at 1%, 5%, and 10% levels, respectively, and the corresponding critical values are 11.344, 7.814, and 6.251 for  $\chi^2(3)$  [Number of Lags = 3], and 23.209, 18.307, and 15.987 for  $\chi^2(10)$  [Number of Lags = 10], and 26.217, 21.026, and 18.549 for  $\chi^2(12)$  [Number of Lags = 12].

The results for different lags  $q$  are reported in Table 9. The latter shows that there is a strong Granger causality from US/UK exchange rate to S&P 500 index price at different expectiles, in particular the lower expectiles and up to 0.7 expectile. We also find some evidence of causality in expectiles from US/Canada and US/Japan exchange rates to S&P 500 index price, although it is less present compared to causality from US/UK exchange rate to S&P 500 index price. The Kolmogorov-Smirnov type-Wald test also confirms the causality from these exchange rates to stock returns, especially when the number lags is high ( $q = 10, 12$ ).

## 6 Conclusion

We proposed a consistent parametric test of Granger causality at a given expectile. We also derived a sup-Wald test for jointly testing Granger causality at all expectiles that has the correct asymptotic size and power properties. Working with expectiles has the advantage of capturing similar information as quantiles, but expectiles has the merit of being much more straightforward to implement than quantiles, since expectiles are define as least squares analogue of quantiles. In other words, studying Granger causality in expectiles is practically simpler and allows us to examine the causality at all levels of the conditional distribution. In addition, testing Granger causality at all expectiles provides a sufficient condition for testing Granger causality in distribution. A Monte Carlo simulation study revealed that our tests have good finite-sample size and power properties for a variety of data-generating processes and different sample sizes. We also compared the properties (size and power) of our expectile-based test with the existing quantile-based test for a range of values of  $\tau$ . The simulation results showed that our test outperforms the test based on quantile regression even for weak degree of causality and in both small and large samples. Finally, we provided two empirical applications to illustrate the usefulness of our tests. In these applications, we re-examined the Granger causality from volume and exchange rates to stock returns using expectile regressions.

## References

- [1] Abdous, B., and B., Remillard. (1995). Relating quantiles and expectiles under weighted-symmetry. *Annals of the Institute of Statistical Mathematics*, 47 (2), 371-384.

- [2] Baur, D. G., T., Dimpfl, and R. C., Jung. (2012). Stock return autocorrelations revisited: A quantile regression approach. *Journal of Empirical Finance*, 19 (2), 254-265.
- [3] Bellini, F., and E. Di., Bernardino. (2017). Risk management with expectiles. *The European Journal of Finance*, 23 (6), 487-506.
- [4] Bellini, F., B. Klar, A. Muller, and E. R., Gianin. (2014). Generalized quantiles as risk measures. *Insurance: Mathematics and Economics*, 54, 41-48.
- [5] Bollerslev, T. (1986). Generalized autoregressive conditional heteroskedasticity. *Journal of Econometrics*, 31 (3), 307-327.
- [6] Boudjellaba, H., J. M., Dufour, and R., Roy. (1992). Testing causality between two vectors in multivariate autoregressive moving average models. *Journal of the American Statistical Association*, 87 (420), 1082-1090.
- [7] Boudjellaba, H., J. M., Dufour, and R., Roy. (1994). Simplified conditions for noncausality between vectors in multivariate ARMA models. *Journal of Econometrics*, 63 (1), 271-287.
- [8] Bouezmarni, T., J. V. Rombouts, and A., Taamouti. (2012). Nonparametric copula-based test for conditional independence with applications to Granger causality. *Journal of Business & Economic Statistics*, 30 (2), 275-287.
- [9] Bouezmarni, T., and A., Taamouti. (2014). Nonparametric tests for conditional independence using conditional distributions. *Journal of Nonparametric Statistics*, 26 (4), 697-719.
- [10] Cenesizoglu, T., and A., Timmermann. (2008). Is the distribution of stock returns predictable? Available at SSRN 1107185.
- [11] Cheung, Y.-W., and L. K., Ng. (1996). A causality-in-variance test and its application to financial market prices. *Journal of Econometrics*, 72 (1-2), 33-48.
- [12] Chow, E.H., W.Y., Lee, and M.E., Solt. (1997). The exchange-rate risk exposure of asset returns. *Journal of Business*, 70,105–123.
- [13] Chuang, C.-C., C.-M. Kuan, and H.-Y., Lin. (2009). Causality in quantiles and dynamic stock return-volume relations. *Journal of Banking & Finance*, 33 (7), 1351-1360.

- [14] Daouia, A., I. Gijbels, and G., Stuper. (2019). Extremiles: A new perspective on asymmetric least squares. *Journal of the American Statistical Association*, 114 (527), 1366-1381.
- [15] Daouia, A., S. Girard, and G., Stuper. (2018). Estimation of tail risk based on extreme expectiles. *Journal of the Royal Statistical Society: Series B*, 80 (2), 263-292.
- [16] Delbaen, F. (2013). A remark on the structure of expectiles. arXiv preprint arXiv:1307.5881.
- [17] Dufour, J.-M., and A., Taamouti. (2010). Short and long run causality measures: Theory and inference. *Journal of Econometrics*, 154 (1), 42-58.
- [18] Efron, B. (1991). Regression percentiles using asymmetric squared error loss. *Statistica Sinica*, 93-125.
- [19] Emmer, S., M. Kratz, and D., Tasche. (2015). What is the best risk measure in practice? a comparison of standard measures. *Journal of Risk*, 18 (2).
- [20] Engle, R. F. (1982). Autoregressive conditional heteroscedasticity with estimates of the variance of United Kingdom inflation. *Econometrica: Journal of the Econometric Society*, 987-1007.
- [21] Fama, E. F., and K. R., French. (1988). Dividend yields and expected stock returns. *Journal of Financial Economics*, 22, 3- 27.
- [22] Granger, C. W. (1969). Investigating causal relations by econometric models and cross-spectral methods. *Econometrica: Journal of the Econometric Society*, 424-438.
- [23] Granger, C. W. (1980). Testing for causality: a personal viewpoint. *Journal of Economic Dynamics and control*, 2, 329-352.
- [24] Granger, C.W.J., and P., Newbold. (1986), *Forecasting Economic Time Series* (2nd ed.), *New York: Academic Press*.
- [25] Granger, C. W. J., R. Robins, and R. F., Engle. (1986). Wholesale and retail prices: Bivariate time series modeling with forecastable error variances. *Model reliability*, 1-17.

- [26] Griffin, J.M., and R.M., Stulz. (2001). International competition and exchange rate shocks: a cross-country industry analysis of stock returns. *Review of Financial Studies*, 14, 215-241.
- [27] Hjort, N. L. and D., Pollard. (1993). Asymptotics for minimisers of convex processes. Technical report, Yale University.
- [28] Holzmann, H., and B., Klar. (2016). Expectile asymptotics. *Electronic Journal of Statistics*, 10 (2), 2355-2371.
- [29] Keim, D. B., and R. F., Stambaugh. (1986). Predicting returns in the stock and bond markets. *Journal of Financial Economics*, 17 (2), 357-390.
- [30] Koenker, R. (2005). Quantile Regression (Econometric Society Monographs; No. 38). *Cambridge university press*.
- [31] Koenker, R., and J. A., Machado. (1999). Goodness-of-fit and related inference processes for quantile regression. *Journal of the American Statistical Association*, 94 (448), 1296-1310.
- [32] Kratschmer, V., and H., Zahle. (2017). Statistical inference for expectile-based risk measures. *Scandinavian Journal of Statistics*, 44 (2), 425-454.
- [33] Kuan, C.-M., J.-H. Yeh, and Y.-C., Hsu. (2009). Assessing value at risk with care, the conditional autoregressive expectile models. *Journal of Econometrics*, 150 (2), 261-270.
- [34] Lee, T.-H. and W., Yang. (2012). Money-income Granger-causality in quantiles. Volume 30, pp. 385-409. *Emerald Group Publishing Limited*.
- [35] Lütkepohl, H. (2005). New introduction to multiple time series analysis. Springer Science & Business Media.
- [36] Newey, W. K., and J. L., Powell. (1987). Asymmetric least squares estimation and testing. *Econometrica: Journal of the Econometric Society*, 819-847.
- [37] Portnoy, S. (1984). Tightness of the sequence of empiric C.D.F processes defined from regression fractiles. In *Robust and Nonlinear Time Series Analysis* (pp. 231-246). Springer, New York, NY.

- [38] Schnabel, S. K., and P. H., Eilers. (2009). Optimal expectile smoothing. *Computational Statistics & Data Analysis*, 53 (12), 4168-4177.
- [39] Sobotka, F., G. Kauermann, L. S. Waltrup, and T., Kneib. (2013). On confidence intervals for semiparametric expectile regression. *Statistics and Computing*, 23 (2), 135-148.
- [40] Sobotka, F. and T., Kneib. (2012). Geoadditive expectile regression. *Computational Statistics & Data Analysis*, 56 (4), 755-767.
- [41] Sobotka, F., S. Schnabel, L. Schulze Waltrup, P. Eilers, T. Kneib, and G., Kauermann. (2011). Expectreg: expectile and quantile regression. R package version 0.25.
- [42] Song, X., and A., Taamouti. (2018). Measuring nonlinear Granger causality in mean. *Journal of Business & Economic Statistics*, 36 (2), 321-333.
- [43] Song, X., and A., Taamouti. (2020). Measuring nonlinear Granger causality in quantiles. *Journal of Business & Economic Statistics* 361-42.
- [44] Su, L., and H., White. (2007). A consistent characteristic function-based test for conditional independence. *Journal of Econometrics*, 141, 807-834.
- [45] Su, L., and H., White. (2008). A nonparametric Hellinger metric test for conditional independence. *Econometric Theory*, 24, 829-864.
- [46] Taamouti, A., T. Bouezmarni, and A., El Ghouch. (2014). Nonparametric estimation and inference for conditional density based Granger causality measures. *Journal of Econometrics*, 180(2), 251-264.
- [47] Taylor, J. W. (2008). Estimating value at risk and expected shortfall using expectiles. *Journal of Financial Econometrics* 6 (2), 231-252.
- [48] Troster, V. (2018). Testing for Granger-causality in quantiles. *Econometric Reviews*, 37 (8), 850-866.
- [49] Waltrup, L. S., F. Sobotka, T. Kneib, and G., Kauermann. (2015). Expectile and quantile regression, david and goliath? *Statistical Modelling*, 15 (5), 433-456.

- [50] Wiener, N. (1956). The theory of prediction. New York: McGraw-Hill.
- [51] Williamson, R. (2001). Exchange rate exposure and competition: evidence from the automotive industry. *Journal of Financial Economics*, 59, 441-475.
- [52] Yang, Z., A. H. Tu, and Y., Zeng. (2014). Dynamic linkages between asian stock prices and exchange rates: new evidence from causality in quantiles. *Applied Economics*, 46 (11), 1184-1201.
- [53] Yao, Q., and H., Tong. (1996). Asymmetric least squares regression estimation: a nonparametric approach. *Journal of Nonparametric Statistics*, 6 (2-3), 273-292.
- [54] Ülkü, N., and O., Onishchenko. (2019). Trading volume and prediction of stock return reversals: Conditioning on investor types' trading. *Journal of Forecasting*, 38 (6), 582-599.
- [55] Ziegel, J. F. (2016). Coherence and elicibility. *Mathematical Finance*, 26 (4), 901-918.

## 7 Appendix A: Proofs

This appendix contains the proof of the main result in the text.

**Poof of Theorem 1:** Consider the check function:

$$\rho(u) = |\tau - \mathbb{I}(u \leq 0)| u^2 = \psi_\tau(u) u^2, \text{ for } \tau \in (0, 1).$$

The derivative of the function  $\rho$  is given by  $\rho'(u) = 2u\psi_\tau(u)$ . The main idea of the proof of Theorem 1 is based on the following decomposition:

$$\rho(u - v) - \rho(u) = -2uv\psi_\tau(u) + v^2\psi_\tau(u) + (2\tau - 1) R(u, v)(u - v), \quad (13)$$

where  $R(u, v) = (\mathbb{I}(u \leq 0) - \mathbb{I}(u \leq v))(u - v) = \int_0^v [\mathbb{I}(u \leq s) - \mathbb{I}(u \leq 0)] ds$ .

Let  $\epsilon_t^\tau = y_t - x'_{t-1}\theta(\tau)$  and  $v_t = \frac{\delta' x_{t-1}}{\sqrt{T}}$ . Denote by

$$H_T(\delta) = \sum_{t=1}^T [\rho(\epsilon_t^\tau - v_t) - \rho(\epsilon_t^\tau)].$$

From (6), we can show that

$$\sqrt{T} \left( \hat{\theta}(\tau) - \theta(\tau) \right) = \arg \min_{\delta \in \mathbb{R}^{d+1}} H_T(\delta).$$

Note that  $H_T(\delta)$  is a convex random objective function that can be decomposed using (13) as follows:

$$H_T(\delta) = H_{1,T}(\delta) + H_{2,T}(\delta) + H_{3,T}(\delta),$$

where

$$\begin{aligned} H_{1,T}(\delta) &= -\delta U_T = \frac{-2}{\sqrt{T}} \sum_{t=1}^T \delta' x_{t-1} \epsilon_t^\tau \psi_\tau(\epsilon_t^\tau), \\ H_{2,T}(\delta) &= \frac{1}{T} \sum_{t=1}^T \delta' x_{t-1} x'_{t-1} \delta \psi_\tau(\epsilon_t^\tau), \\ H_{3,T}(\delta) &= (2\tau - 1) \sum_{t=1}^T R(\epsilon_t^\tau, v_t) (\epsilon_t^\tau - v_t). \end{aligned}$$

Now, first using the fact that  $0 \leq R(\epsilon^\tau, v)$  and that  $|R(\epsilon^\tau, v) (\epsilon^\tau - v)| = R(\epsilon^\tau, v) |(\epsilon^\tau - v)| \leq R(\epsilon^\tau, v) |v|$ , we obtain

$$|H_{3,T}(\delta)| \leq |(2\tau - 1)| \frac{|\delta' x_{t-1}|}{\sqrt{T}} \sum_{t=1}^T R(\epsilon_t^\tau, v_t).$$

From Koenker(2005, pages 121-122), we have  $\sum_{t=1}^T R(\epsilon_t^\tau, v_t) = O_p(1)$ . Hence, one can deduces that  $H_{2,T}(\delta) = o_p(1)$ . Second, under Assumptions **A.1** and **A.2** and the fact that  $\Psi_\tau = E[\psi_\tau(\epsilon_t^\tau) x_{t-1} x'_{t-1}]$  is finite, we have:

$$H_{2,T}(\delta) = \frac{1}{T} \sum_{t=1}^T \delta' x_{t-1} x'_{t-1} \delta \psi_\tau(\epsilon_t^\tau) = \delta' E[x_{t-1} x'_{t-1} \psi_\tau(\epsilon_t^\tau)] \delta + o_p(1) = \delta' \Psi_\tau \delta + o_p(1).$$

Third, since  $x_{t-1} \in I_{YZ}(t-1)$ , then  $E[\epsilon_t^\tau \psi_\tau(\epsilon_t^\tau) | I_{YZ}(t-1)] = 0$  and  $x_{t-1} \epsilon_t^\tau \psi_\tau(\epsilon_t^\tau)$  is a martingale difference sequence. Hence  $H_{1,T}(\delta)$  satisfies the conditions of central limit theorem. Following similar arguments in Portnoy (1984), we can show the tightness of the autoregressive and thus the weak convergence of the empirical process  $\frac{1}{\sqrt{T}} \sum_{t=1}^T x_{t-1} \epsilon_t^\tau \psi_\tau(\epsilon_t^\tau)$ , i.e.,

$$\frac{1}{\sqrt{T}} \sum_{t=1}^T x_{t-1} \epsilon_t^\tau \psi_\tau(\epsilon_t^\tau) \Longrightarrow \mathcal{Z}(\tau),$$

where  $\mathcal{Z}(\tau)$  is a Gaussian process with covariance function

$$Cov(\mathcal{Z}(\tau_1), \mathcal{Z}(\tau_2)) = \Psi_{\tau_1}^{-1} \Psi_{\tau_1, \tau_2} \Psi_{\tau_2}^{-1},$$

where  $\Psi_{\tau_i} = E[\psi_{\tau_i}(\epsilon_t^{\tau_i}) x_{t-1} x'_{t-1}]$ , for  $i = 1, 2$ , and  $\Psi_{\tau_1, \tau_2} = E[\psi_{\tau_1}(\epsilon_t^{\tau_1}) \psi_{\tau_2}(\epsilon_t^{\tau_2}) \epsilon_t^{\tau_1} \epsilon_t^{\tau_2} x_{t-1} x'_{t-1}]$ .

Finally, from Assumption **A.3**, the basic corollary of Hjort and Pollard (1993), and

$$H_T = \frac{-2}{\sqrt{T}} \sum_{t=1}^T \delta' x_{t-1} \epsilon_t^\tau \psi_\tau(\epsilon_t^\tau) + \frac{1}{T} \sum_{t=1}^T \delta' x_{t-1} x'_{t-1} \delta \psi_\tau(\epsilon_t^\tau) + o_p(1) \xrightarrow{P} -2\delta' \mathcal{Z}(\tau) + \delta' \Psi_\tau \delta,$$

we can deduce that

$$\sqrt{T} \left( \hat{\theta}(\tau) - \theta(\tau) \right) \Longrightarrow -\Psi_\tau^{-1} \mathcal{Z}(\tau).$$

Which concludes the proof of Theorem 1.