

THE UNIVERSITY
of LIVERPOOL

SECTION A

1. Write down the Taylor series about $x = 3$ for the function

$$f(x) = x^{-1}.$$

State whether this Taylor series converges to $f(x)$ for:

- a) $x = 0$, b) $x = 4$. [5 marks]

2. Find the solutions of the following differential equations:

- (i) $e^y \frac{dy}{dx} = x^2$ with $y(1) = 0$,
(ii) $x \frac{dy}{dx} + 2y = 3$ with $y(1) = 0$.

[8 marks]

3. Solve the differential equation

$$\frac{d^2y}{dx^2} + 4\frac{dy}{dx} + 3y = 0$$

with the initial conditions $y(0) = 2$, $y'(0) = 1$.

[5 marks]

4. Show, by taking limits along two different paths to the origin $(0, 0)$, that

$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy + 2x^2}{x^2 + xy + y^2}$$

does not exist.

[4 marks]

THE UNIVERSITY
of LIVERPOOL

5. Work out all first and second partial derivatives of

$$f(x, y) = 4x^3y - 4xy^3,$$

and verify that

$$\frac{\partial^2 f}{\partial x \partial y} = \frac{\partial^2 f}{\partial y \partial x},$$
$$\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} = 0.$$

[5 marks]

6. Suppose that $x = x(t)$, $y = y(t)$, and $z = z(t)$ are functions of t such that

$$x(0) = 2, \quad y(0) = -1, \quad z(0) = 0.$$

Suppose that the derivatives satisfy

$$x'(0) = y'(0) = 1, \quad z'(0) = -1.$$

Then work out

$$\frac{dF}{dt}(0)$$

where $F(t) = f(x(t), y(t), z(t))$, and

$$f(x, y, z) = y^2z + \cos(xyz).$$

[5 marks]

7. Find the gradient vector $\nabla f(1, 1, 1)$, where

$$f(x, y, z) = \frac{x + y + z}{x^2 + y^2 + z^2}.$$

Find also the derivative of f in the direction $\mathbf{i} - 2\mathbf{j} - 2\mathbf{k}$ at $(1, 1, 1)$. [5 marks]

8. Locate and classify all stationary points of the function

$$f(x, y) = x^2 + xy^2 - 4xy - 5x.$$

[8 marks]

THE UNIVERSITY
of LIVERPOOL

9. Find the linear approximation near $(x, y) = (1, 1)$ to the function

$$f(x, y) = \frac{1}{2x^2 - y^2}.$$

[4 marks]

10. By changing the order of integration, compute

$$\int_0^1 \int_y^1 \cos(y/x) dx dy.$$

[6 marks]

SECTION B

11.

In this question, let

$$f(y) = (1 - y)^{-1/2}, \quad g(x) = (1 - x^2)^{-1/2}, \quad h(x) = \sin^{-1}(x).$$

(i) Compute the first five derivatives of f . Write down the 4'th Taylor polynomial of f at 0. Hence, or otherwise, compute the 9th and 10th Taylor polynomials respectively of g and h , at 0.

(ii) Show that if $0 \leq y \leq \frac{1}{4}$, the 4th remainder term $R_4(y, 0)$ of f at 0 satisfies

$$|R_4(y, 0)| \leq \frac{63}{256} \cdot \frac{4^{11/2}}{3^{11/2}} y^5.$$

[15 marks]

12. Solve the following differential equations with the given boundary conditions:

(i)

$$y'' + 4y' - 5y = 4e^{-x}$$

with $y(0) = 1$, $y'(0) = -1$.

(ii)

$$y'' + 4y' - 5y = -5x^2 + 3x + 1$$

with $y(0) = 1$, $y'(0) = -1$.

[15 marks]

13. Find the maximum and minimum values of the function $f(x, y)$ in the region where $g(x, y) \leq 3$, where $f(x, y)$ and $g(x, y)$ are defined by

$$f(x, y) = xy - y,$$
$$g(x, y) = x^2 + 3y^2.$$

[15 marks]

14.

a) Find the weight of the parallelogram R bounded by $y = x$, $x = 0$, $y = x + 1$ and $x = 1$, where the density function is $\rho(x, y) = x$

b) Find the centre of mass (\bar{x}, \bar{y}) of R .

[15 marks]