


# Gaia




## Contents

Why map the stars in the Milky Way?

How can they be mapped accurately?

What type of data can be acquired?

What can we learn from the data?



UNIVERSITY OF LIVERPOOL

2

A slide titled 'Contents' with four bullet points: 'Why map the stars in the Milky Way?', 'How can they be mapped accurately?', 'What type of data can be acquired?', and 'What can we learn from the data?'. Below the text is an image of the Gaia spacecraft. At the bottom left is the University of Liverpool logo, and at the bottom right is the number '2'.


## Mapping the Stars

It's a simple question ...

*Where are the stars?*

We want to be able to determine the position (on the sky) of any celestial object so that we can ...

- ... find it again at a later time
- ... tell if it has moved
- ... tell if any new objects appear



3


A slide titled 'Mapping the Stars'. It contains a list of questions: 'It's a simple question ...', 'Where are the stars?', 'We want to be able to determine the position (on the sky) of any celestial object so that we can ...', and three bullet points: '... find it again at a later time', '... tell if it has moved', and '... tell if any new objects appear'. To the left of the text is a star map of the Serpens Caput region, showing various stars and constellations. At the bottom left is the number '3'.

## Mapping the Stars

It's a simple question ...

*Where are the stars?*

Also, if we can determine the positions of stars in our galaxy with high precision, then we can gain a better understanding of the structure and the history of the Milky Way.



4

A slide titled 'Mapping the Stars'. It contains a list of questions: 'It's a simple question ...', 'Where are the stars?', and 'Also, if we can determine the positions of stars in our galaxy with high precision, then we can gain a better understanding of the structure and the history of the Milky Way.' To the left of the text is a star map of the Serpens Caput region, showing various stars and constellations. At the bottom left is the number '4'.

## Mapping the Stars

So how do we map the sky? Can we just take lots of photos of the night sky and stitch them together?

That approach seems to work fine for mapping the surface of the Earth from space, so what's the problem?

**Problem #1 – Stars move**

( nothing in the galaxy is static )

## Mapping the Stars

**Problem #2 – Stars are at different distances**

Knowing the positions of stars on the 2-dimensional sky is not enough to determine how they are arranged in 3-dimensions.



We also need to know their distances.


## Mapping the Stars

Being able to determine the distances to stars is important not only for understanding the structure of the Milky Way ...

... but also for understanding the stars themselves.

Without knowing the distance to a star we cannot determine its luminosity or its type.

Without its type, we cannot determine the properties of any of its exoplanets.




## Parallax

Parallax is the apparent change in the position of a celestial object due to the motion of the Earth around the Sun.


Diagrams like this always imply that a (nearby) star can be aligned with more distant (fixed) stars to determine the parallax angle.

However, in practice, *all* stars appear to move as the Earth orbits the Sun.



## Parallax

Parallax is the apparent change in the position of a celestial object due to the motion of the Earth around the Sun.



A star ~3 ly distant would show a parallax angle of ~1 arc second.

This distance is defined as one "parallax second".

1 parsec = 3.26 light-years


## Measuring Parallax

So here is the crux of the problem.


- To calculate distances we need parallax
- Parallax needs the stars' *actual* positions in the sky (not simply their position *relative* to other stars)
- The stars themselves are moving (proper motion)

This is why mapping the stars in the Milky Way is not a trivial task.

How can they be mapped accurately?



## Mapping the Stars



### Measuring Parallax

To untangle the apparent motion of a star due to parallax from its proper motion as it moves within the Milky Way we need

- High precision measurements <<< parallax angles are **very small**
- A fixed coordinate system <<< Earth **and** stars are in motion

[ Aside – There were plans in the 1980s to make the parallax angles larger (and so easier to measure) by increasing the baseline from 1AU to 1000AU. Flying a spacecraft with an ion drive that far out into the solar system would take 50 years. For a sense of scale, the Voyager spacecraft are currently ~ 150AU from Earth. ]



### Coordinate Systems

What coordinate system should we use to map the stars?


The equatorial coordinate system, comprising Right Ascension (RA) and Declination (Dec) coordinates, uses the celestial equator as a reference.

RA and Dec have been used in star atlases since the days of the first Astronomer Royal.


Why are they not good enough?



### Coordinate Systems



Epoch 1950.0




Epoch 2000.0



### Coordinates Relative to the Sun

So if an Earth-based coordinate system is not suitable, what about using the Sun as a reference? That would also lead to small errors because the Sun orbits around the solar system's centre of mass.




We think of planets orbiting the Sun, but actually they all orbit around a common **barycentre**


Hence the best coordinate system uses this barycentre as a fixed reference point.

# Gaia

## Parallax Only




## Parallax and Proper Motion



## Hipparcos

High Precision PARallax COLlecting Satellite 1989–1993

Hipparcos was the first space telescope dedicated to high-precision measurements of star positions. Its name was also a homage to the Greek astronomer Hipparchus, the founder of trigonometry.



Hipparcos catalogued 100,000 stars with a precision of  $\sim 0.001$  arcsec.

(  $0.001$  arcsec = milli arcsec = mas )

Faintest stars = mag 12


The Hipparcos catalogue was used by the Hubble Space Telescope.

## Gaia

Global Astrometric Interferometer for Astrophysics 2013–2025

( Note that the acronym is no longer relevant as its design changed )

Gaia's mission was to improve on Hipparcos by mapping a billion stars with a precision of  $\sim 10 \mu\text{as}$ .  
(  $1 \mu\text{as} = 0.001$  mas )



Higher precision meant that stars at greater distances could have their parallax measured, and so Gaia could 'reach' further into the Milky Way.

Faintest stars = mag 20

# Gaia

## Micro Arc Seconds

An analogy for Gaia's ability to measure tiny angles precisely...

$10 \mu\text{as} = 0.01 \text{ mas} = 0.00001 \text{ arc seconds}$

= the thickness of a human hair  
seen from a distance of 1000 km




## Alternatives?

Why build a new telescope to achieve this? Why not use ...




Hubble  
Space Telescope



Simonyi (aka LSST)  
in Vera Rubin Obs

|                              |          |           |
|------------------------------|----------|-----------|
| Precision                    | < 1 mas  | 100 mas   |
| # stars                      | 1000     | > billion |
| Time to do<br>all-sky survey | > 1000 y | 10 y      |

How does Gaia  
achieve this?




## Gaia Hardware



Gaia was ... constructed 2006–2012

... launched in Dec 2013

... placed into a halo orbit  
around L2 in Jan 2014



# Gaia

## Why Park at L2?

At L2 a sunshield can block light from the Sun, Earth and Moon

UNIVERSITY OF LIVERPOOL

25

## Sunshield Deployment

UNIVERSITY OF LIVERPOOL

26

## Gaia Hardware

Gaia was ... constructed 2006–2012

... launched in Dec 2013

... placed into a halo orbit around L2 in Jan 2014

How was Gaia's micro arcsec ( $\mu\text{as}$ ) precision achieved?

Would it need a big mirror, a few km in diameter? Obviously not.


UNIVERSITY OF LIVERPOOL

27

Space Telescopes

UNIVERSITY OF LIVERPOOL

# Gaia





## Gaia Hardware

Gaia achieves its high precision through an ingenious combination of instrumental hardware and a novel mode of operating:

### Hardware

- A large (Gigapixel) detector array comprising 106 CCD sensors
- Two telescopes that simultaneously image different patches of the sky





## Gaia Operation

Gaia achieves its high precision through an ingenious combination of instrumental hardware and a novel mode of operating:

### Operation


- Both telescopes create star images on the **same** detector array
- Gaia spins and so the star images **drift** across the detector CCDs
- Gaia does **not** take images



## Gaia Operation

This makes Gaia very different to imaging space telescopes like the Hubble or James Webb Space Telescopes.


Reading data from the CCDs is a crucial element of the operation and so it is worth taking a closer look at how this is done.



000001010  
011100101  
110111000  
000001010  
011100101  
110111000

# Gaia

## CCD Images



Imagine an image of a star field.


How is the image made?

Image data is just an array of numbers.

UNIVERSITY OF LIVERPOOL

33

## Reading a CCD In 'Stare' Mode



Expose CCD to light

Exposure complete


Read data from CCD

Read complete

UNIVERSITY OF LIVERPOOL

34

## Reading a CCD In 'Stare' Mode



Expose CCD to light

Exposure complete

Read data from CCD


Read complete

1001001011  
0111

UNIVERSITY OF LIVERPOOL

35

## Reading a CCD In 'Stare' Mode



Expose CCD to light

Exposure complete

Read data from CCD

Read complete


1001001011  
0111100100  
0101101110  
1011100010  
1001001011

UNIVERSITY OF LIVERPOOL

36


# Gaia

## Reading a CCD In 'Drift' Mode




As Gaia spins, stars drift across the CCDs.

## Reading a CCD In 'Drift' Mode



Stars drift across CCD  
CCD pixels shift in synch  
Signal keeps adding up  
Star reaches readout  
Pixels are read out

## Reading a CCD In 'Drift' Mode



Stars drift across CCD  
CCD pixels shift in synch  
Signal keeps adding up  
Star reaches readout  
Pixels are read out

Star




## Two Telescopes, One Detector

A sextant works on a similar principle ...







Light from two objects is reflected from mirrors so that they appear together in the 'same detector'.







For navigation, the altitude of the star (red) is used to determine the observer's latitude.





# Gaia







# Gaia



# Gaia




# Gaia




# Gaia

DR2

All-Sky Map



What can we learn  
from the data?




DR2

H-R Diagram

Hertzsprung–Russell diagrams plot the luminosity (vertical) against the colour (horizontal) for a population of stars.





This H-R diagram for millions of stars in the Milky Way gives us a way to visualise the stellar population of our galaxy.







DR3

Sun-like Star and Neutron Star









DR3


## Sagittarius Dwarf Galaxy Mergers



## Star Streams







## Star Streams



## Star Streams



# Gaia





# Gaia

## Gaia Epilogue

As it drifted away from L2 it was bright enough to be imaged by amateur astrophotographers.

Its heliocentric orbit will bring it close to Earth every 14 years.


Before it was powered down Gaia sent a final status update.

UNIVERSITY OF LIVERPOOL

77

## Gaia Epilogue



NGR


HST

Field of view of the Nancy Grace Roman Space Telescope

UNIVERSITY OF LIVERPOOL

78

## Gaia Epilogue



Milky Way Dynamics and Evolution

Dark Matter

Dark Energy

Solar System

Exoplanets

Neutron Stars

Black Holes


Stellar Evolution

DR4 = 500 TB

UNIVERSITY OF LIVERPOOL

79

[www.liverpool.ac.uk/~sdb/Talks](http://www.liverpool.ac.uk/~sdb/Talks)



# Gaia

Dr Steve Barrett

BADAS 8 Jul 2026