



Contents

Why map the stars in the Milky Way?

How can they be mapped accurately?


What type of data can be acquired?

What can we learn from the data?



UNIVERSITY OF LIVERPOOL

2



Mapping the Stars

It's a simple question…

Where are the stars?


We want to be able to determine the position (on the sky) of any celestial object so that we can…

…find it again at a later time

…tell if it has moved

…tell if any new objects appear

3



Mapping the Stars

It's a simple question…

Where are the stars?

Also, if we can determine the positions of stars in our galaxy with high precision, then we can gain a better understanding of the structure and the history of the Milky Way.

4

Mapping the Stars

So how do we map the sky? Can we just take lots of photos of the night sky and stitch them together?

That approach seems to work fine for mapping the surface of the Earth from space, so what's the problem?

Problem #1 – Stars move

(nothing in the galaxy is static)

UNIVERSITY OF LIVERPOOL

Mapping the Stars

Problem #2 – Stars are at different distances

Knowing the positions of stars on the 2-dimensional sky is not enough to determine how they are arranged in 3-dimensions.



We also need to know their distances.

UNIVERSITY OF LIVERPOOL


Mapping the Stars

Being able to determine the distances to stars is important not only for understanding the structure of the Milky Way ...

... but also for understanding the stars themselves.

Without knowing the distance to a star we cannot determine its luminosity or its type.

Without its type, we cannot determine the properties of any of its exoplanets.




UNIVERSITY OF LIVERPOOL

Parallax

Parallax is the apparent change in the position of a celestial object due to the motion of the Earth around the Sun.

Diagrams like this always imply that a (nearby) star can be aligned with more distant (fixed) stars to determine the parallax angle.


However, in practice, ***all*** stars appear to move as the Earth orbits the Sun.



UNIVERSITY OF LIVERPOOL

Parallax

Parallax is the apparent change in the position of a celestial object due to the motion of the Earth around the Sun.



A star ~3 ly distant would show a parallax angle of ~1 arc second.

This distance is defined as one "parallax second".

1 parsec = 3.26 light-years


Measuring Parallax

So here is the crux of the problem.


- To calculate distances we need parallax
- Parallax needs the stars' *actual* positions in the sky (not simply their position *relative* to other stars)
- The stars themselves are moving (proper motion)

This is why mapping the stars in the Milky Way is not a trivial task.

How can they be mapped accurately?



Mapping the Stars



Measuring Parallax

To untangle the apparent motion of a star due to parallax from its proper motion as it moves within the Milky Way we need

- High precision measurements <<< parallax angles are ***very*** small
- A fixed coordinate system <<< Earth ***and*** stars are in motion

[Aside – There were plans in the 1980s to make the parallax angles larger (and so easier to measure) by increasing the baseline from 1 AU to 1000 AU. Flying a spacecraft with an ion drive that far out into the solar system would take 50 years. For a sense of scale, the Voyager spacecraft are currently ~ 150 AU from Earth.]

UNIVERSITY OF LIVERPOOL


Coordinate Systems

What coordinate system should we use to map the stars?


The equatorial coordinate system, comprising Right Ascension (RA) and Declination (Dec) coordinates, uses the celestial equator as a reference.

RA and Dec have been used in star atlases since the days of the first Astronomer Royal.


Why are they not good enough?




UNIVERSITY OF LIVERPOOL



Coordinate Systems



Epoch 1950.0




Epoch 2000.0



Coordinates Relative to the Sun

So if an Earth-based coordinate system is not suitable, what about using the Sun as a reference? That would also lead to small errors because the Sun orbits around the solar system's centre of mass.



We think of planets orbiting the Sun, but actually they all orbit around a common *barycentre*

Hence the best coordinate system uses this barycentre as a fixed reference point.


UNIVERSITY OF LIVERPOOL

Parallax Only



UNIVERSITY OF LIVERPOOL 17

Parallax and Proper Motion




UNIVERSITY OF LIVERPOOL 18

Hipparcos

HIgh **P**recision **PAR**allax **CO**llecting **S**atellite 1989 – 1993

Hipparcos was the first space telescope dedicated to high-precision measurements of star positions. Its name was also a homage to the Greek astronomer Hipparchus, the founder of trigonometry.



Hipparcos catalogued 100,000 stars with a precision of ~0.001 arcsec.

(0.001 arcsec = milli arcsec = mas)

Faintest stars = mag 12

The Hipparcos catalogue was used by the Hubble Space Telescope.


UNIVERSITY OF LIVERPOOL 19

Gaia

Global **A**strometric **I**nterferometer for **A**strophysics 2013 – 2025

(Note that the acronym is no longer relevant as its design changed)

Gaia's mission was to improve on Hipparcos by mapping a billion stars with a precision of ~10 μas. (1 μas = 0.001 mas)



Higher precision meant that stars at greater distances could have their parallax measured, and so Gaia could 'reach' further into the Milky Way.

Faintest stars = mag 20

UNIVERSITY OF LIVERPOOL 20

Micro Arc Seconds

An analogy for Gaia's ability to measure tiny angles precisely…

10 μas = 0.01 mas = 0.00001 arc seconds

= the thickness of a human hair
seen from a distance of 1000 km



UNIVERSITY OF LIVERPOOL


21

Alternatives?

Why build a new telescope to achieve this? Why not use…



Hubble
Space Telescope




Simonyi (aka LSST)
in Vera Rubin Obs

	Hubble Space Telescope	Simonyi (aka LSST) in Vera Rubin Obs
Precision	< 1 mas	100 mas
# stars	1000	> billion
Time to do all-sky survey	> 1000 y	10 y

UNIVERSITY OF LIVERPOOL

22

How does Gaia achieve this?



UNIVERSITY OF LIVERPOOL


Gaia Hardware

esa ASTRIUM AIRBUS

Gaia was … constructed 2006–2012

… launched in Dec 2013


… placed into a halo orbit
around L2 in Jan 2014



UNIVERSITY OF LIVERPOOL

24

Why Park at L2?



UNIVERSITY OF LIVERPOOL

25

Sunshield Deployment



UNIVERSITY OF LIVERPOOL

26


Gaia Hardware

esa ASTRIUM AIRBUS

Gaia was ... constructed 2006–2012

... launched in Dec 2013

... placed into a halo orbit around L2 in Jan 2014





How was Gaia's micro arcsec (μas) precision achieved?

Would it need a big mirror, a few km in diameter? Obviously not.

UNIVERSITY OF LIVERPOOL

27







Gaia Hardware

Gaia achieves its high precision through an ingenious combination of instrumental hardware and a novel mode of operating:

Hardware

- A large (Gigapixel) detector array comprising 106 CCD sensors
- Two telescopes that simultaneously image different patches of the sky





UNIVERSITY OF LIVERPOOL


30


Gaia Operation

Gaia achieves its high precision through an ingenious combination of instrumental hardware and a novel mode of operating:

Operation

- Both telescopes create star images on the ***same*** detector array
- Gaia spins and so the star images ***drift*** across the detector CCDs
- Gaia does ***not*** take images






UNIVERSITY OF LIVERPOOL

31

Gaia Operation

This makes Gaia very different to imaging space telescopes like the Hubble or James Webb Space Telescopes.

Reading data from the CCDs is a crucial element of the operation and so it is worth taking a closer look at how this is done.




```
000001010
011100101
110111000
000001010
011100101
110111000
```

UNIVERSITY OF LIVERPOOL

32

CCD Images



Imagine an image of a star field.


How is the image made?

Image data is just an array of numbers.

UNIVERSITY OF LIVERPOOL

33

Reading a CCD In 'Stare' Mode



Expose CCD to light

Exposure complete


Read data from CCD

Read complete

UNIVERSITY OF LIVERPOOL

34

Reading a CCD In 'Stare' Mode



Expose CCD to light

Exposure complete

Read data from CCD


Read complete

1001001011
0111

UNIVERSITY OF LIVERPOOL

35

Reading a CCD In 'Stare' Mode



Expose CCD to light

Exposure complete

Read data from CCD


Read complete

1001001011
0111100100
0101101110
1011100010
1001001011

UNIVERSITY OF LIVERPOOL


36

Reading a CCD In 'Drift' Mode



As Gaia spins, stars drift across the CCDs.

Reading a CCD In 'Drift' Mode



Stars drift across CCD


CCD pixels shift in synch

Signal keeps adding up

Star reaches readout


Pixels are read out

Reading a CCD In 'Drift' Mode



Two Telescopes, One Detector


A sextant works on a similar principle ...



For navigation, the altitude of the star (red) is used to determine the observer's latitude.

Light from two objects is reflected from mirrors so that they appear together in the 'same detector'.


Gaia Optical Bench



UNIVERSITY OF LIVERPOOL

41


Gigapixel CCD Array



UNIVERSITY OF LIVERPOOL


42

Operation of Gaia



UNIVERSITY OF LIVERPOOL

Gaia Operation




Two telescopes but one detector

UNIVERSITY OF LIVERPOOL

44

Stars Drift Across CCD




Star identified by sky mapper

UNIVERSITY OF LIVERPOOL

45

Stars Drift Across CCD




Star transits timed by astrometrics CCDs

UNIVERSITY OF LIVERPOOL

46

Stars Drift Across CCD




Star brightness measured in blue and red

UNIVERSITY OF LIVERPOOL

47

Stars Drift Across CCD




Star spectrum recorded

UNIVERSITY OF LIVERPOOL


48

Stars Drift Across CCD




Thousands of stars measured per second

Gaia Operation



Gaia spins once every 6 hours

Its spin axis precesses around the Gaia-Sun axis every 63 days



It covers the whole sky in under a year

Data Releases DR1–2

2014 2016 2018 2020 2022 2024 2026 2028 2030

Original Mission | Extended Mission

Full data

Data Release 1 2 E 3 F 4 5

Coords of ...
~ millions
~ billions
of stars

Data Release DR3

2014 2016 2018 2020 2022 2024 2026 2028 2030

Original Mission | Extended Mission

Full data

Data Release 1 2 E 3 F 4 5


Coords of ...
~ millions
~ billions
of stars

Improved precision


~ millions of variable stars
~ millions of galaxies
~ 100 million spectra
~ 100 thousand SS objects

Gaia


Data Release FPR




Milky Way Spiral Arms




Region of Galaxy Covered



DR3 Region Covered By Gaia




DR2 **All-Sky Map**



UNIVERSITY OF LIVERPOOL

57

What can we learn from the data?




UNIVERSITY OF LIVERPOOL

DR2 **H-R Diagram**

Hertzsprung–Russell diagrams plot the luminosity (vertical) against the colour (horizontal) for a population of stars.


This H-R diagram for millions of stars in the Milky Way gives us a way to visualise the stellar population of our galaxy.



UNIVERSITY OF LIVERPOOL

59


DR3 **Sun-like Star and Neutron Star**



UNIVERSITY OF LIVERPOOL


60

DR3 **Hidden Neutron Stars**




UNIVERSITY OF LIVERPOOL 61

DR2 **Warped Disc**




UNIVERSITY OF LIVERPOOL 62

Warped Disc



UNIVERSITY OF LIVERPOOL 63

DR3 **Stellar Motion**



UNIVERSITY OF LIVERPOOL 64

Galaxy Mergers

Hierarchical model of galaxy formation




UNIVERSITY OF LIVERPOOL

65

DR2+DR3

Galaxy Merger History




UNIVERSITY OF LIVERPOOL

66

DR3

Sagittarius Dwarf Galaxy Mergers




UNIVERSITY OF LIVERPOOL

67

DR3

Sagittarius Dwarf Galaxy Mergers




UNIVERSITY OF LIVERPOOL

68

DR3

Sagittarius Dwarf Galaxy Mergers



UNIVERSITY OF LIVERPOOL

69


Star Streams



UNIVERSITY OF LIVERPOOL

70

Star Streams



UNIVERSITY OF LIVERPOOL

71

Star Streams




UNIVERSITY OF LIVERPOOL

72

DR2+DR3

Galaxy Merger History




UNIVERSITY OF LIVERPOOL

73

DR2+DR3


Galaxy Merger Future



UNIVERSITY OF LIVERPOOL


74

Epilogue



UNIVERSITY OF LIVERPOOL

Gaia Epilogue




UNIVERSITY OF LIVERPOOL

76

Gaia Epilogue

As it drifted away from L2 it was bright enough to be imaged by amateur astrophotographers.



Its heliocentric orbit will bring it close to Earth every 14 years.


Before it was powered down Gaia sent a final status update.



UNIVERSITY OF LIVERPOOL

77


Gaia Epilogue




UNIVERSITY OF LIVERPOOL

78

Gaia Epilogue





DR4 = 500 TB

Milky Way
Dynamics and Evolution


Dark Matter
Dark Energy
Solar System
Exoplanets
Neutron Stars
Black Holes
Stellar Evolution

UNIVERSITY OF LIVERPOOL

79

www.liverpool.ac.uk/~sdb/Talks

Gaia



Dr Steve Barrett

SU3A 23 Nov 2026