It's Not Rocket Science!



It's Not Rocket Science!

Well, actually, it is…
…but it's not THAT hard
if you just skip the maths

Dr Steve Barrett
UKStratCom
29 Mar 2023

It's Not Rocket Science

UNIVERSITY OF LIVERPOOL 2

Contents

UNIVERSITY OF LIVERPOOL 3

What Does a Rocket Do?

Q: What is the primary function of a rocket?

A: To lift objects up above the atmosphere.

Well, yes … sort of.



The ISS and the HST are actually still **in** the atmosphere – it's just rather thin up there at an altitude of ~500 km.

To put an object into Earth orbit what is needed is horizontal speed.

(Getting as high as possible to reduce atmospheric drag is a good idea, but it is not strictly necessary.)

UNIVERSITY OF LIVERPOOL 4

It's Not Rocket Science!

Newton's Cannonballs



Isaac Newton
Principia 1686



The need for horizontal velocity was realised more than 300 years ago

UNIVERSITY OF LIVERPOOL

Saturn V Launch

UNIVERSITY OF LIVERPOOL

Saturn V Launch

UNIVERSITY OF LIVERPOOL

Otherwise: What Goes Up …



UNIVERSITY OF LIVERPOOL

Rocket Science
What does a rocket do?

Orbits
Kepler, Newton and Buzz Aldrin
Circular – elliptical – parabolic – hyperbolic

Getting From A to B
Earth to Mars
Gravity assists to the rest of the solar system

Parking Places
What are Lagrange points all about?

Orbits

Orbit : The path of an object affected by (only) gravity.

We are all used to the idea of planets in orbit around the Sun, or moons in orbit around their planets.



Kepler observed that the planets orbit in ellipses and Newton figured out why – his law of universal gravitation.

Conic Sections

Newton found that orbits have the shapes of sections through a cone



Conic Sections

These curves that describe orbits around the Sun are also the curves that define the shapes of mirrors that are suitable for focusing light.

This is because the maths is basically the same for both.



Planets follow closed elliptical orbits, but a comet falling into the solar system from way out in the Oort cloud follows a parabolic path.

Objects entering and leaving the solar system at speed follow a hyperbolic orbit.

It's Not Rocket Science!

Hyperbolic Orbit



UNIVERSITY OF LIVERPOOL

13

Orbit Size and Orbit Period

Green planet is closer to Sun

Gravitational pull from Sun is stronger

Planet moves faster to stay in orbit

Shorter period ('year')



Red planet is further from Sun

Gravitational pull from Sun is weaker

Planet moves more slowly to stay in orbit

Longer period ('year')

UNIVERSITY OF LIVERPOOL

14

Earth Satellite Orbits



UNIVERSITY OF LIVERPOOL

15

Planets Orbit the Sun



UNIVERSITY OF LIVERPOOL

16

Circular Orbits

Two objects with equal mass orbit around the centre of mass (+)

If the objects have different masses, objects with larger mass move smaller distances



If the masses are very different, the centre of mass (+) may lie within the larger object

Even a small object (planet) can make a large object (star) wobble – this is how exoplanets can be detected

UNIVERSITY OF LIVERPOOL 17

3-Body Problem

Describing the relative motion of two objects is a soluble problem



but for three or more objects there are very few analytic solutions

UNIVERSITY OF LIVERPOOL 18

Rocket Science

What does a rocket do?

Orbits

Kepler, Newton and Buzz Aldrin

Circular – elliptical – parabolic – hyperbolic

Getting From A to B

Earth to Mars

Gravity assists to the rest of the solar system

Parking Places

What are Lagrange points all about?

UNIVERSITY OF LIVERPOOL 19

Earth to Mars



UNIVERSITY OF LIVERPOOL 20

Hohmann Transfer



Not to scale

Hohmann Transfer



Not to scale

Hohmann Transfer

… and another to get into the new orbit.



A Hohmann transfer needs a kick to get out of Earth's orbit…

Gravity Assists

What about getting to other planets in the solar system?

In the 1960s it was realised that flying a spacecraft close to a planet can 'slingshot' it onwards at higher velocities.



Hence exploring the outer solar system can be carried out faster and cheaper.

In effect, the spacecraft robs the planet of some of its momentum.


Falling Into a Gravity Well
Same height
Same speed
Max speed
UNIVERSITY OF
LIVERPOOL
25


Jupiter Gravity Assist
Inward
speed
Outward
speed
Max flyby speed
UNIVERSITY OF
LIVERPOOL
26


Jupiter Gravity Assist
Inward
speed
Jupiter
orbits the Sun
Outward
speed
10 km/s
Relative to the Sun
25 km/s
Relative to the Sun
Max flyby speed
UNIVERSITY OF
LIVERPOOL
27


Pioneer 10
1972-03-03
Pioneer 10
Earth
Jupiter
0.0km/s
831,951,662km
UNIVERSITY OF
LIVERPOOL
28

Pioneer 11



Voyager 1



Voyager 2



Voyager 2 Speed vs Distance



It's Not Rocket Science!

Gravity Assists

Using gravity assists to help cover the enormous distances between planets in the outer solar system seems like a very sensible idea.

What is not so obvious is that gravity assists are also used to visit the ***inner*** planets.

A planet flyby can be used to ***lose*** speed as a spacecraft 'falls' into the inner solar system.



A flyby can accelerate ***or*** decelerate, but in both cases fuel is saved.

Changing Orbits Uses Fuel

Some spacecraft go into orbit around their target planet



Any orbital changes require fuel which reduces the mission lifetime

Rocket Science
- What does a rocket do?

Orbits
- Kepler, Newton and Buzz Aldrin
- Circular – elliptical – parabolic – hyperbolic

Getting From A to B
- Earth to Mars
- Gravity assists to the rest of the solar system

Parking Places
- What are Lagrange points all about?

Orbit Size and Orbit Period

Green planet is closer to Sun

Gravitational pull from Sun is stronger

Planet moves faster to stay in orbit

Shorter period ('year')



Red planet is further from Sun

Gravitational pull from Sun is weaker

Planet moves more slowly to stay in orbit

Longer period ('year')

Orbit Size and Orbit Period

The arrows show the gravitational force of the Sun on each planet.

At a greater distance, the force is less.

Is that always the case?



We can't change the gravitational pull of the Sun…

…but we can arrange it so that an object in the red orbit feels an additional gravitational force.

Lagrange Points

Let's assume that green is planet Earth with an orbital period of exactly 1 year.

The red orbit has a period of ***more*** than 1 year.

What if the red orbit is close enough to the Earth that an object is pulled by both the Sun ***AND*** the Earth?



There is a red orbit at *just* the right distance such that the extra pull from the Earth makes the red object orbit faster with a period of 1 year.

That would mean that the red object would orbit the Sun 'with' the Earth.

Lagrange Point L2



Lagrange Point L2



Not to scale

It's Not Rocket Science!

Lagrange Point L1

The idea is conceptually the same for L1 *inside* the Earth's orbit, but this time the Sun and the Earth pull in opposite directions.



UNIVERSITY OF LIVERPOOL 41

Lagrange Point L1

L1 is a good location for spacecraft that observe the Sun, as the Earth never gets in the way.



UNIVERSITY OF LIVERPOOL 42

Calculating Lagrange Points

Gravitational forces between two bodies fall off as the square of the distance between them, so we can use that to calculate the distance from Earth to L1 or L2:

$$\frac{M}{(R \pm r)^2} \pm \frac{m}{r^2} = \left(\left(\frac{M}{M+m}\right)R \pm r\right)\left(\frac{M+m}{R^3}\right)$$

M = mass of Sun R = distance Sun – Earth
m = mass of Earth r = distance Earth – L1/2

Just solve for r. Simples!

UNIVERSITY OF LIVERPOOL 43

Calculating Lagrange Points

Gravitational forces between two bodies fall off as the square of the distance between them, so we can use that to calculate the distance from Earth to L1 or L2:

$$\frac{M}{(R \pm r)^2} \pm \frac{m}{r^2} = \left(\left(\frac{M}{M+m}\right)R \pm r\right)\left(\frac{M+m}{R^3}\right)$$

M = mass of Sun R = distance Sun – Earth
m = mass of Earth r = distance Earth – L1/2

Just solve for r. Simples!

... but it's not THAT hard if you just skip the maths

UNIVERSITY OF LIVERPOOL 44


Lagrange Points 1 – 5
L4
L3
L1
L2
L5
UNIVERSITY OF LIVERPOOL
45


Why Park at L2?
L4
L3
L1
At L2 a sunshield can block light from the Sun, Earth and Moon
L5
UNIVERSITY OF LIVERPOOL
46


Why Park at L2?
L2 is in Earth's shadow, so spacecraft like Gaia and JWST orbit around L2
L4
L3
L1
If a spacecraft moves away from L2 then the pull of the Earth brings it back
L5
UNIVERSITY OF LIVERPOOL
47


Why Park at L2?
L2 is in Earth's shadow, so spacecraft like Gaia and JWST orbit around L2
L4
L3
L1
If a spacecraft moves towards or away from Earth then it will continue to drift
L5
UNIVERSITY OF LIVERPOOL
48

It's Not Rocket Science!

Parking at L2



The ability of a spacecraft to 'park' at a Lagrange point is sometimes visualised using this doughnut.

Imagine that the contours represent height and think about a ball wanting to roll 'downhill'.

Look more closely at the contours around L2 …

UNIVERSITY OF LIVERPOOL

Parking at L2



The arrows show the directions of 'downhill' in the vicinity of L2.

L2 is *not* on a 'hilltop'.

L2 is *not* in a 'well'.

L2 is at a saddle point.



UNIVERSITY OF LIVERPOOL

Parking at L2



Hence the JWST can park in a 'halo' orbit around L2, but it will slowly drift towards Earth …



… and so will need a station-keeping nudge every few months.

UNIVERSITY OF LIVERPOOL

Objects at L4 and L5



UNIVERSITY OF LIVERPOOL

Objects at L4 and L5



Objects can orbit loosely around the L4 and L5 points.

There's not much there at the Earth L4 and L5 points.

UNIVERSITY OF LIVERPOOL

53

Objects at L4 and L5



Objects can orbit loosely around the L4 and L5 points.

There's not much there at the Earth L4 and L5 points.

Jupiter's stronger gravitational pull has collected some Trojan asteroids at its L4 and L5 points.

UNIVERSITY OF LIVERPOOL

54

Horseshoe Orbits



Some orbits look really weird …

… but they still obey the rules:

When closer to the Sun the object goes faster than the Earth.

When further from the Sun the object goes slower than the Earth.

UNIVERSITY OF LIVERPOOL

55

Open Orbits

Perhaps the weirdest of orbits are those that are ***not*** closed loops



Is it possible for the stars in a triple-star system to move this way?

UNIVERSITY OF LIVERPOOL

56

Open Orbits

...and don't get me started on orbits around black holes



UNIVERSITY OF LIVERPOOL 

57

Summary

Rocket Science
What does a rocket do?

Orbits
Kepler, Newton and Buzz Aldrin
Circular – elliptical – parabolic – hyperbolic

Getting From A to B
Earth to Mars
Gravity assists to the rest of the solar system

Parking Places
What are Lagrange points all about?

UNIVERSITY OF LIVERPOOL

58

Summary

Why is "rocket science" considered to be so hard?



Yes, there are some objects in the solar system that move in ways that are not, at first sight, intuitive.

Yes, calculating how to put the JWST into orbit around L2 was not trivial (especially given the accuracy achieved).



But the underlying idea is straightforward enough: stronger gravitational pull leads to faster motion. (The Earth moves around the Sun faster than Mars)

That's really all there is to it. The rest is ***just*** maths.

After all ... it's not rocket science!

UNIVERSITY OF LIVERPOOL

59