



Contents

[A Brief History](#)

[Science Fact](#)

[Science Fiction](#)

[The Future](#)



2

A Brief History

Black Hole [definition]:

A region of space from which nothing, not even light, can escape.


When did the concept of a black hole originate? Earlier than you think.

Rev John Michell (1783) – "If [the size of] a sphere of the same density of the Sun were to exceed that of the Sun in the proportion of 500 to 1 ... all light emitted from such a body would be made to return towards it by its gravity."

In the 1800s the idea of such '**dark stars**' was largely ignored.




3



Warping Space and Time


A Brief History

	Einstein (1915) General Theory of Relativity		Schwarzschild (1916) Calculation of gravity of a compact mass
	Eddington (1926) Stars compressed to the 'Schwarzschild radius'		Chandrasekhar (1931) Massive stars at the end of their lives will collapse
	Oppenheimer (1939) Nothing can stop the collapse of massive stars	General Relativity predicts that time stops at the Schwarzschild radius and so such collapsed stars were called ' frozen stars '.	

UNIVERSITY OF LIVERPOOL

5

A Brief History

Finkelstein (1958)	
At the Schwarzschild radius there is an event horizon – a 'one-way membrane'	
Golden Age of GR and BH (~ 1960 – 1975)	
Kerr / Newman / Penrose / Hawking	
1964 First use of the term ' black hole '	
1967 Discovery of pulsars	
1969 Identified as neutron stars	
Gravitational collapse of stars is not just a hypothetical possibility!	

UNIVERSITY OF LIVERPOOL

6

Pieces of the Puzzle

UNIVERSITY OF LIVERPOOL

7

UNIVERSITY OF LIVERPOOL

8

Warping Space and Time




Escape Velocity

What goes up must come down? Right?

Wrong, not if it goes up fast enough.

The threshold speed needed for an object to escape the gravitational pull of a body is called the **escape velocity**. If you throw something up with a speed less than the escape velocity then it will fall back.

The force of gravity pulling an object (like you) towards a body (like the Earth) depends on both of the masses involved and the distance between them. The escape velocity does not depend on the mass of the object — the escape velocity for a 1 kg satellite is the same as for a 1 ton satellite or even a 1000 ton space station.



UNIVERSITY OF
LIVERPOOL

9

10

The escape velocity at the surface of a body depends on the ratio of the mass to the radius

$$\frac{M}{R}$$

Some escape velocities:

Velocity to escape the **Moon** = 2 km/s (= 5,000 mph)

Velocity to escape the **Earth** = 11 km/s (= 25,000 mph)

Velocity to escape the **Sun** = 600 km/s (> 1,000,000 mph)

600 km/s may sound like a lot, but it is only 0.2% of the speed of light




UNIVERSITY OF
LIVERPOOL



Escape Velocity

Let's compress the Sun to smaller sizes and see what happens to the escape velocity.



UNIVERSITY OF
LIVERPOOL

11

12



Escape Velocity

If we compress the Sun into a sphere of radius 3 km then we form a black hole.

What if we start with a body with a different mass?

For the **Earth** the Schwarzschild radius is ~ cm

Moon

~ 0.1 mm

Mt Everest



We know of no way that this can happen for any mass smaller than that of a star, but that doesn't mean that it's impossible.



UNIVERSITY OF
LIVERPOOL

12


Warping Space and Time



Escape Velocity

Although all sizes are possible, small BH don't last.

Aside: Quantum Mechanics* allows particles and antiparticles to be created from borrowed energy, as long as they annihilate and pay back the borrowed energy on very short time scales.



UNIVERSITY OF
LIVERPOOL

* See "The Weird World of the Very Very Small"


13



Escape Velocity

Although all sizes are possible, small BH don't last.

How is this particle-antiparticle creation relevant to the lifetime of BH? What might happen if they are created *just outside* the event horizon?



There is a net flux of particles radiating from the event horizon of the BH called **Hawking radiation**. This radiation increases with decreasing mass, so small BH evaporate! More about this later in the talk.



UNIVERSITY OF
LIVERPOOL

14



Escape Velocity

What happens if we compress an object to a size *smaller* than the Schwarzschild radius?

The event horizon, the point at which the escape velocity is equal to the speed of light, is still the same (the Schwarzschild radius). →




For massive stars ($M > 10 M_{\odot}$) nothing that we know of can stop the core collapse during a supernova and so all the mass is compressed to an infinitely small point – a **singularity**.

We cannot tell if this is what actually happens inside a BH because it is all hidden inside the event horizon. Could we ever see a "naked" singularity? Not according to the **Cosmic Censorship Hypothesis**.



UNIVERSITY OF
LIVERPOOL

15



General Relativity



UNIVERSITY OF
LIVERPOOL

16

Warping Space and Time



General Relativity

Einstein's General Theory of Relativity
First presented in 1915 and published in 1916

What does it say?

$$G = 8\pi T$$

The main principle of GR can be expressed by this one equation that looks deceptively simple.

G is the geometry of space
 T is the distribution of mass

Mass distorts space!



UNIVERSITY OF
LIVERPOOL

17



General Relativity

$$G = 8\pi T$$

The equation is sometimes reduced to the more prosaic description

Mass tells space how to curve

Space tells mass how to move

What do we mean by curved space?

Any image trying to explain this uses the analogy of 2-dimensional space curving into a third dimension.

To warp or curve 3-dimensional space we need a fourth dimension which humans, not unnaturally, find very difficult to imagine.




UNIVERSITY OF
LIVERPOOL

18



General Relativity

Black holes produce infinitely warped space




UNIVERSITY OF
LIVERPOOL

19




Predictions of General Relativity

The precession of the orbit of Mercury



Gravitational lensing



Stars seen during the total solar eclipse of 1919 were found to be slightly shifted in position.



UNIVERSITY OF
LIVERPOOL

20

Warping Space and Time

 **Predictions of General Relativity**

Time slows down in a gravitational field





UNIVERSITY OF LIVERPOOL

21

 **Gravitational Waves**

One of the predictions of GR is gravitational waves.



The ripples in spacetime are tiny — 1 part in 10^{20}

UNIVERSITY OF LIVERPOOL

22

 **Gravitational Waves**

Merger of two neutron stars




UNIVERSITY OF LIVERPOOL

23

 **Gravitational Waves**

If binary pulsars emit gravitational waves they must be slowly losing energy, and this would mean that they would be expected to slow down (black line). This is precisely what is observed (blue points).



Year	Period shift in seconds
1970	0
1975	-1
1980	-2
1985	-4
1990	-7
1995	-12
2000	-25

UNIVERSITY OF LIVERPOOL

24

Warping Space and Time



A black hole, represented by a small black circle, is shown interacting with a 2x2 grid of four puzzle pieces. The pieces are colored purple, green, yellow, and red. The black hole is positioned above the yellow piece, which is partially overlapping the red piece.

Stars and Galaxies

UNIVERSITY OF LIVERPOOL

25




Black Holes Are Born

Core collapse of a massive star – Supernova*

UNIVERSITY OF LIVERPOOL

* See "The ABC of Stellar Evolution"

26




Black Holes Feed

Infalling matter forms an accretion disk

UNIVERSITY OF LIVERPOOL

27



Black Holes Feed

Accretion disk and x-ray jets

Accretion disks form because objects passing close to the BH are ripped apart by huge tidal forces.

The effect is more popularly known as spaghettiification or the noodle effect.

UNIVERSITY OF LIVERPOOL

28

Warping Space and Time



Black Holes Feed

Accretion disk and x-ray jets



The infalling matter is heated by the tidal forces and emits light and x-rays.

The accretion process can convert more than 25% of the infalling mass into energy.

Compare this to nuclear fusion in stars, where less than 1% of the mass is converted into energy.



29



Black Holes Feed


Is this how black holes feed on neighbouring stars?



30



Black Hole Jets



HERCULES A

HTTP://CHANDRA.GSFC.NASA.GOV




31



Supermassive Black Holes

Zooming in to the centre of the Milky Way




32

Warping Space and Time

 **Supermassive Black Holes**

Stars orbiting the BH at the centre of the Milky Way




UNIVERSITY OF LIVERPOOL

33

 **Supermassive Black Holes**

How big is the SMBH at the centre of the Milky Way?

From the orbits of the stars the SMBH mass is determined to be $4 \text{ million } M_{\odot}$




Other SMBH are much larger:

Andromeda Galaxy	$250 \text{ million } M_{\odot}$
Sombrero Galaxy	$1 \text{ billion } M_{\odot}$
Largest SMBH	$> 20 \text{ billion } M_{\odot}$

UNIVERSITY OF LIVERPOOL

34




Detecting Gravity

UNIVERSITY OF LIVERPOOL

35

 **Measuring Curved Space**

Gravity Probe B
2004



UNIVERSITY OF LIVERPOOL

36

Warping Space and Time

Detecting Extreme Gravity

SWIFT
Gamma-Ray Burst Mission
2004

FERMI
Gamma-Ray Space Telescope
2008

UNIVERSITY OF LIVERPOOL

37

This slide shows two space telescopes: SWIFT (launched 2004) and FERMI (launched 2008). The SWIFT satellite is shown in an orange and yellow color scheme, while the FERMI satellite is blue. Both have solar panels deployed. The University of Liverpool logo is at the bottom left.

Detecting Extreme Gravity

The gamma-ray cycle of PG 1553+113

Gamma-ray flux

2009 2010 2011 2012 2013 2014 2015

UNIVERSITY OF LIVERPOOL

38

This slide features a composite image showing a gamma-ray burst source and a plot titled "The gamma-ray cycle of PG 1553+113". The plot shows the gamma-ray flux over time from 2009 to 2015, with data points represented by purple diamonds and error bars. The University of Liverpool logo is at the bottom left.

Detecting Gravity Waves

LIGO
Laser
Interferometer
Gravitational
Wave
Observatory

UNIVERSITY OF LIVERPOOL

39

This slide shows the LIGO Laser Interferometer Gravitational Wave Observatory. It features a large aerial photograph of the LIGO facility in Louisiana, with a red box highlighting a specific part of the interferometer. A small inset image shows a top-down view of the interferometer's internal structure. The University of Liverpool logo is at the bottom left.

Detecting Gravity Waves


UNIVERSITY OF LIVERPOOL

40

This slide provides a detailed look inside the LIGO interferometer. It shows the massive vacuum tanks and the complex optical equipment used to detect gravitational waves. The University of Liverpool logo is at the bottom left.


Warping Space and Time

 **Detecting Gravity Waves**




UNIVERSITY OF LIVERPOOL

41



 **Detecting Gravity Waves**

eLISA = Evolved Laser Interferometer Space Antenna



UNIVERSITY OF LIVERPOOL

43

 **Detecting Gravity Waves**

eLISA = Evolved Laser Interferometer Space Antenna

LIGO is sensitive to waves with periods of less than a second. This means that it can detect waves created by the merger of two black holes, but this does not happen very often.



By comparison, eLISA will be sensitive to waves with much longer periods, from seconds to hours. This should mean that eLISA can detect signals from BH within our galaxy and SMBH in other galaxies.



It is also hoped that eLISA will be able to detect the gravitational waves created by the biggest singularity of them all – the Big Bang.*

* See "The Beginning of Everything"

UNIVERSITY OF LIVERPOOL

44

Warping Space and Time



Measuring an Event Horizon


Event Horizon Telescope



UNIVERSITY OF
LIVERPOOL

45

Escape Velocity General Relativity



We have looked at the science facts

Now let's look at the science fiction

Stars and Galaxies

Detecting Gravity




UNIVERSITY OF
LIVERPOOL

46

Pieces of the Puzzle

Does Hollywood ever get it right?



Ship is parked right next to the event horizon



Ship is pulled in to a planetary black hole



Ship is powered by an onboard black hole



UNIVERSITY OF
LIVERPOOL

47



If the radiation from such a BH, smaller than an atom, could be captured or directed, then that is enough energy to power a starship.



UNIVERSITY OF
LIVERPOOL

48

Science Fiction

Is a BH-powered star drive possible?
For BH with mass **less** than a kg, not much energy is available ($E=mc^2$).

For BH with a mass **more** than that of Mount Everest, Hawking radiation is very weak.

For BH with mass ~ 1 million tons the power in the Hawking radiation would be equivalent to the output of ~ 100 million nuclear power stations and the BH would last ~ 100 years.


Warping Space and Time

Science Fiction

What about shortcuts through space?

Wormholes?

Hyperspace?



49

Science Fiction

What about shortcuts through space?

Wormholes?

Hyperspace?



50

Science Fiction

What about time travel?



51

Science Fiction

Hawking proposed the
Chronology Protection Conjecture

A wormhole allowing time travel will
collapse before anything has time to
travel through it.

"Making the Universe safe for historians"

What about time travel?



52

Warping Space and Time

The Problem

There's a problem.

General Relativity (GR) works really well for **massive** objects (like stars).

Quantum Mechanics (QM) works really well for **tiny** objects (like atoms).*

But, what if the object is **massive and tiny**?

Then we need (drum roll)... **Quantum Gravity**.

The problem: GR and QM are not good bedfellows.

A universal 'Theory of Everything' has proven to be elusive.



* See "The Weird World of the Very Very Small"

53



Maybe quantum mechanics will prevent a singularity from forming, thus avoiding the horrible properties like infinite density and infinitely warped space.

For instance, String Theory describes a ten-dimensional universe in which the fundamental building blocks are 'strings' rather than the more familiar 'particles'.



If String Theory is right, black holes are 'fuzzballs' without a singularity at their core. They are just 'balls of string'.

But, is the universe described by String Theory the one in which we live?




54

The Future

Einstein never really believed that quantum mechanics was the right description of the microscopic world.

He spent most of his later years wrestling with a Theory of Everything.

If a genius like Einstein could not get his head around the problem, what will it take?



Maybe some unexpected discoveries, for instance from LIGO or eLISA, will point the way forward to a better understanding of black holes.



55

Warping Space and Time

www.liverpool.ac.uk/~sdb/Talks

Dr Steve Barrett

King's School 25 Mar 2019