

Warping Space and Time



Contents

A Brief History

Science Fact

Science Fiction

The Future

A Brief History

Black Hole [definition]:

A region of space from which nothing, not even light, can escape.

When did the concept of a black hole originate? Earlier than you think.

Rev John Michell (1783) – "If [the size of] a sphere of the same density of the Sun were to exceed that of the Sun in the proportion of 500 to 1 ... all light emitted from such a body would be made to return towards it by its gravity."

In the 1800s the idea of such '**dark stars**' was largely ignored.


A Brief History




Einstein (1915)
General Theory of Relativity



Schwarzschild (1916)
Calculation of gravity
of a compact mass



Eddington (1926)
Stars compressed to the
'Schwarzschild radius'



Chandrasekhar (1931)
Massive stars at the end
of their lives will collapse



Oppenheimer (1939)
Nothing can stop the
collapse of massive stars

General Relativity predicts that time stops at the Schwarzschild radius and so such collapsed stars were called '**frozen stars**'.

Warping Space and Time

A Brief History

Finkelstein (1958)
At the Schwarzschild radius there is an event horizon – a 'one-way membrane'


Golden Age of GR and BH (~ 1960–1975)
Kerr / Newman / Penrose / Hawking

1964 First use of the term '**black hole**'

1967 Discovery of pulsars

1969 Identified as neutron stars

Gravitational collapse of stars is not just a hypothetical possibility!



Pieces of the Puzzle

Escape
Velocity


General
Relativity

We will look at each of these
pieces of the puzzle in turn

Stars and
Galaxies

Detecting
Gravity

Escape
Velocity



Escape Velocity




What goes up must come down? Right?

Wrong, not if it goes up fast enough.

The threshold speed needed for an object to escape the gravitational pull of a body is called the **escape velocity**. If you throw something up with a speed less than the escape velocity then it will fall back.

The force of gravity pulling an object (like you) towards a body (like the Earth) depends on both of the masses involved and the distance between them. The escape velocity does not depend on the mass of the object — the escape velocity for a 1 kg satellite is the same as for a 1 ton satellite or even a 1000 ton space station.



Warping Space and Time

Escape Velocity

The escape velocity at the surface of a body depends on the ratio of the mass to the radius

$$\frac{M}{R}$$


Some escape velocities:

Velocity to escape the **Moon** = 2 km/s (= 5,000 mph)
 Velocity to escape the **Earth** = 11 km/s (= 25,000 mph)
 Velocity to escape the **Sun** = 600 km/s (> 1,000,000 mph)

600 km/s may sound like a lot, but it is only 0.2% of the speed of light

UNIVERSITY OF LIVERPOOL

9



Escape Velocity

If we compress the Sun into a sphere of radius 3 km then we form a black hole.

What if we start with a body with a different mass?

For the **Earth** the Schwarzschild radius is ~ cm
Moon ~ 0.1 mm
Mt Everest ~ atom

We know of no way that this can happen for any mass smaller than that of a star, but that doesn't mean that it's impossible.

UNIVERSITY OF LIVERPOOL

11

Escape Velocity

Although all sizes are possible, small BH don't last.

Aside: Quantum Mechanics* allows particles and antiparticles to be created from borrowed energy, as long as they annihilate and pay back the borrowed energy on very short time scales.

UNIVERSITY OF LIVERPOOL

* See "The Weird World of the Very Very Small"


12

Warping Space and Time

Escape Velocity

Although all sizes are possible, small BH don't last.

How is this particle-antiparticle creation relevant to the lifetime of BH?
What might happen if they are created *just* outside the event horizon?



There is a net flux of particles radiating from the event horizon of the BH called **Hawking radiation**. This radiation increases with decreasing mass, so small BH evaporate! More about this later in the talk.


UNIVERSITY OF LIVERPOOL

13

Escape Velocity

What happens if we compress an object to a size *smaller* than the Schwarzschild radius?

The event horizon, the point at which the escape velocity is equal to the speed of light, is still the same (the Schwarzschild radius).




For massive stars ($M > 10M_{\odot}$) nothing that we know of can stop the core collapse during a supernova and so all the mass is compressed to an infinitely small point – a **singularity**.

We cannot tell if this is what actually happens inside a BH because it is all hidden inside the event horizon. Could we ever see a "naked" singularity? Not according to the **Cosmic Censorship Hypothesis**.

UNIVERSITY OF LIVERPOOL

14

General Relativity



UNIVERSITY OF LIVERPOOL

15

General Relativity


Einstein's General Theory of Relativity
First presented in 1915 and published in 1916

What does it say? $G = 8\pi T$

The main principle of GR can be expressed by this one equation that looks deceptively simple.

G is the geometry of space
 T is the distribution of mass


Mass distorts space!



UNIVERSITY OF LIVERPOOL

16

Warping Space and Time




General Relativity

$$G = 8\pi T$$


The equation is sometimes reduced to the more prosaic description

Mass tells **space** how to **curve**
Space tells **mass** how to **move**

What do we mean by curved space?
Any image trying to explain this uses the analogy of 2-dimensional space curving into a third dimension.
To warp or curve 3-dimensional space we need a fourth dimension which humans, not unnaturally, find very difficult to imagine.





17




General Relativity

Black holes produce infinitely warped space







18




Predictions of General Relativity




The precession of the orbit of Mercury




Gravitational lensing



Stars seen during the total solar eclipse of 1919 were found to be slightly shifted in position.







19



Predictions of General Relativity


Time slows down in a gravitational field






20


Warping Space and Time


 **Gravitational Waves**


One of the predictions of GR is gravitational waves.





The ripples in spacetime are tiny — 1 part in 10^{20}


 21




 **Stars and Galaxies**




 23


 **Black Holes Are Born**

Core collapse of a massive star – Supernova*





 * See "The ABC of Stellar Evolution" 24


Warping Space and Time

 **Black Holes Feed**


Infalling matter forms an accretion disk



 25


 **Black Holes Feed**


Accretion disk and x-ray jets




Accretion disks form because objects passing close to the BH are ripped apart by huge tidal forces.

The effect is more popularly known as spaghettification or the noodle effect.

 26

 **Black Holes Feed**


Accretion disk and x-ray jets



The infalling matter is heated by the tidal forces and emits light and x-rays.

The accretion process can convert more than 25% of the infalling mass into energy.

Compare this to nuclear fusion in stars, where less than 1% of the mass is converted into energy.

 27


 **Black Holes Feed**

Is this how black holes feed on neighbouring stars?




 28

Warping Space and Time


 **Black Hole Jets**

HERCULES A




UNIVERSITY OF LIVERPOOL

29


 **Supermassive Black Holes**

Stars orbiting the BH at the centre of the Milky Way




UNIVERSITY OF LIVERPOOL

30

 **Supermassive Black Holes**

How big is the SMBH at the centre of the Milky Way?


From the orbits of the stars the SMBH mass is determined to be 4 million M_{\odot}



Other SMBH are much larger:


Andromeda Galaxy	250 million M_{\odot}
Sombrero Galaxy	1 billion M_{\odot}
Largest SMBH	> 20 billion M_{\odot}

UNIVERSITY OF LIVERPOOL

 **Black Hole In Star Cluster**

Using the same technique, the Hubble Space Telescope has detected a massive black hole at the core of this star cluster.


Do all large clusters contain a black hole?



UNIVERSITY OF LIVERPOOL

32


Warping Space and Time



Detecting Gravity


UNIVERSITY OF LIVERPOOL

33




Measuring Curved Space

Gravity Probe B
2004




UNIVERSITY OF LIVERPOOL

34




Detecting Extreme Gravity

SWIFT
Gamma-Ray Burst Mission
2004




FERMI
Gamma-Ray Space Telescope
2008




UNIVERSITY OF LIVERPOOL

35



Detecting Extreme Gravity



The gamma-ray cycle of PG 1553+113

Gamma-ray flux
Extraterrestrial


2009 2010 2011 2012 2013 2014 2015

UNIVERSITY OF LIVERPOOL

36

Warping Space and Time

Detecting Gravitational Waves




LIGO
Laser
Interferometer
Gravitational
Wave
Observatory

UNIVERSITY OF LIVERPOOL

37


Detecting Gravitational Waves



UNIVERSITY OF LIVERPOOL


38

Detecting Gravitational Waves




UNIVERSITY OF LIVERPOOL


39





Warping Space and Time

 **Detecting Gravitational Waves**

eLISA = Evolved Laser Interferometer Space Antenna




 41


 **Detecting Gravitational Waves**

eLISA = Evolved Laser Interferometer Space Antenna


LIGO is sensitive to waves with periods of less than a second. This means that it can detect waves created by the merger of two black holes, but this does not happen very often.



By comparison, eLISA will be sensitive to waves with much longer periods, from seconds to hours. This should mean that eLISA can detect signals from BH within our galaxy and SMBH in other galaxies.



It is also hoped that eLISA will be able to detect the gravitational waves created by the biggest singularity of them all – the Big Bang.*


 * See "The Beginning of Everything" 42


 **Event Horizon Telescope**




M87*

 43

 **Event Horizon Telescope**




Sgr A*

 44

Warping Space and Time

Event Horizon Telescope



M87*

Sgr A*

Voyager 1

Pluto's orbit


Sun's diameter

Mercury's orbit

UNIVERSITY OF LIVERPOOL

45

Pieces of the Puzzle



Escape Velocity

General Relativity

We have looked at the science facts

Now let's look at the science fiction

Stars and Galaxies

Detecting Gravity

UNIVERSITY OF LIVERPOOL

46

Science Fiction

Does Hollywood ever get it right?



Ship is parked right next to the event horizon


Ship is pulled in to a planetary black hole

Ship is powered by an onboard black hole

UNIVERSITY OF LIVERPOOL

47

Science Fiction



Is a BH-powered star drive possible?

For BH with mass **less** than a kg, not much energy is available ($E = mc^2$).

For BH with a mass **more** than that of Mount Everest, Hawking radiation is very weak.

For BH with mass ~ 1 million tons the power in the Hawking radiation would be equivalent to the output of ~ 100 million nuclear power stations and the BH would last ~ 100 years.

If the radiation from such a BH, smaller than an atom, could be captured or directed, then that is enough energy to power a starship.

UNIVERSITY OF LIVERPOOL

48


Warping Space and Time

Science Fiction

What about shortcuts through space?

Wormholes?

Hyperspace?



Science Fiction

What about shortcuts through space?

Wormholes?

Hyperspace?



Science Fiction

What about time travel?



Science Fiction

What about time travel?

Hawking proposed the

Chronology Protection Conjecture

A wormhole allowing time travel will collapse before anything has time to travel through it.

"Making the Universe safe for historians"



Warping Space and Time

The Problem

There's a problem.

General Relativity (GR) works really well for **massive** objects (like stars).

Quantum Mechanics (QM) works really well for **tiny** objects (like atoms).*

But, what if the object is massive **and** tiny?

Then we need (drum roll)... **Quantum Gravity**.

The problem: GR and QM are not good bedfellows.

A universal 'Theory of Everything' has proven to be elusive.



* See "The Weird World of the Very Very Small"

53

The Future



Maybe quantum mechanics will prevent a singularity from forming, thus avoiding the horrible properties like infinite density and infinitely warped space.

For instance, String Theory describes a ten-dimensional universe in which the fundamental building blocks are 'strings' rather than the more familiar 'particles'.

If String Theory is right, black holes are 'fuzzballs' without a singularity at their core. They are just 'balls of string'.

But, is the universe described by String Theory the one in which we live?




54

The Future

Einstein never really believed that quantum mechanics was the right description of the microscopic world.

He spent most of his later years wrestling with a Theory of Everything.

If a genius like Einstein could not get his head around the problem, what will it take?



Maybe some unexpected discoveries, for instance from LIGO or eLISA, will point the way forward to a better understanding of black holes.



55

Warping Space and Time

www.liverpool.ac.uk/~sdb/Talks

Dr Steve Barrett

MAC 10 Oct 2023